

表 2587 体重(kg) ヘモグロビン濃度(g/dL)別（腹膜透析患者）

ヘモグロビン濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
5未満			1		1	3	1	1					7		7	62.44	12.79
(%)			(14.3)		(14.3)	(42.9)	(14.3)	(14.3)					(100.0)				
5～				1	5	5	2	1					14		14	61.87	11.24
(%)				(7.1)	(35.7)	(35.7)	(14.3)	(7.1)					(100.0)				
6～			1	3	8	7	4		1				24	4	28	60.11	12.40
(%)			(4.2)	(12.5)	(33.3)	(29.2)	(16.7)		(4.2)				(100.0)				
7～	1	2	2	21	33	26	19	4	4				112	12	124	60.22	15.02
(%)	(0.9)	(1.8)	(1.8)	(18.8)	(29.5)	(23.2)	(17.0)	(3.6)	(3.6)				(100.0)				
8～	4	4	12	47	78	71	46	21	7	3		2	295	49	344	60.93	16.18
(%)	(1.4)	(1.4)	(4.1)	(15.9)	(26.4)	(24.1)	(15.6)	(7.1)	(2.4)	(1.0)		(0.7)	(100.0)				
9～	7		30	127	232	229	117	43	23	2	2	1	813	90	903	60.99	14.15
(%)	(0.9)		(3.7)	(15.6)	(28.5)	(28.2)	(14.4)	(5.3)	(2.8)	(0.2)	(0.2)	(0.1)	(100.0)				
10～	14	9	41	246	423	456	218	86	18	7	3	2	1,523	183	1,706	60.68	13.88
(%)	(0.9)	(0.6)	(2.7)	(16.2)	(27.8)	(29.9)	(14.3)	(5.6)	(1.2)	(0.5)	(0.2)	(0.1)	(100.0)				
11～	7	4	44	249	461	469	230	70	23	2	1	2	1,562	147	1,709	60.47	13.00
(%)	(0.4)	(0.3)	(2.8)	(15.9)	(29.5)	(30.0)	(14.7)	(4.5)	(1.5)	(0.1)	(0.1)	(0.1)	(100.0)				
12～	3	2	26	129	210	213	100	35	14	3	2	1	738	58	796	60.38	13.71
(%)	(0.4)	(0.3)	(3.5)	(17.5)	(28.5)	(28.9)	(13.6)	(4.7)	(1.9)	(0.4)	(0.3)	(0.1)	(100.0)				
13～	1	1	3	33	60	55	38	14	7	2	1	1	216	21	237	63.24	15.99
(%)	(0.5)	(0.5)	(1.4)	(15.3)	(27.8)	(25.5)	(17.6)	(6.5)	(3.2)	(0.9)	(0.5)	(0.5)	(100.0)				
14～	1	1	5	11	20	20	8	7	1				74	5	79	59.04	14.88
(%)	(1.4)	(1.4)	(6.8)	(14.9)	(27.0)	(27.0)	(10.8)	(9.5)	(1.4)				(100.0)				
合計	38	23	165	867	1,531	1,554	783	282	98	19	9	9	5,378	569	5,947	60.71	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.1)	(28.5)	(28.9)	(14.6)	(5.2)	(1.8)	(0.4)	(0.2)	(0.2)	(100.0)				
記載なし			7	37	60	62	28	16	6	1	1		218	2,776	2,994	61.51	13.75
(%)			(3.2)	(17.0)	(27.5)	(28.4)	(12.8)	(7.3)	(2.8)	(0.5)	(0.5)		(100.0)				
総計	38	23	172	904	1,591	1,616	811	298	104	20	10	9	5,596	3,345	8,941	60.74	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.2)	(28.4)	(28.9)	(14.5)	(5.3)	(1.9)	(0.4)	(0.2)	(0.2)	(100.0)				
平均	10.48	10.43	10.81	10.86	10.83	10.86	10.81	10.79	10.69	10.78	10.97	10.68	10.83	10.64	10.81		
標準偏差	1.43	1.74	1.50	1.39	1.40	1.41	1.43	1.52	1.59	1.44	1.44	1.83	1.42	1.37	1.42		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査