

表 2590 体重(kg) Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
50未満				3	7	4	2	2					18		18	61.14	13.20
(%)				(16.7)	(38.9)	(22.2)	(11.1)	(11.1)					(100.0)				
50～			3	10	26	29	22	8	4				102	12	114	64.32	13.26
(%)			(2.9)	(9.8)	(25.5)	(28.4)	(21.6)	(7.8)	(3.9)				(100.0)				
70～			9	47	76	103	61	19	6	1	2	1	325	26	351	63.15	13.46
(%)			(2.8)	(14.5)	(23.4)	(31.7)	(18.8)	(5.8)	(1.8)	(0.3)	(0.6)	(0.3)	(100.0)				
90～			17	70	176	176	106	48	17		1	1	612	48	660	63.20	13.43
(%)			(2.8)	(11.4)	(28.8)	(28.8)	(17.3)	(7.8)	(2.8)		(0.2)	(0.2)	(100.0)				
110～			17	127	194	235	99	35	11	3	1	1	723	74	797	61.14	12.53
(%)			(2.4)	(17.6)	(26.8)	(32.5)	(13.7)	(4.8)	(1.5)	(0.4)	(0.1)	(0.1)	(100.0)				
130～	1	2	12	103	186	177	80	24	15	3		2	605	57	662	60.97	13.48
(%)	(0.2)	(0.3)	(2.0)	(17.0)	(30.7)	(29.3)	(13.2)	(4.0)	(2.5)	(0.5)		(0.3)	(100.0)				
150～	2	2	10	71	116	128	51	20	7			2	409	27	436	60.18	13.63
(%)	(0.5)	(0.5)	(2.4)	(17.4)	(28.4)	(31.3)	(12.5)	(4.9)	(1.7)			(0.5)	(100.0)				
170～			6	54	66	59	34	21	4				244	19	263	60.35	13.04
(%)			(2.5)	(22.1)	(27.0)	(24.2)	(13.9)	(8.6)	(1.6)				(100.0)				
190～	1		3	18	38	25	9	9	1	2		1	107	13	120	60.25	15.60
(%)	(0.9)		(2.8)	(16.8)	(35.5)	(23.4)	(8.4)	(8.4)	(0.9)	(1.9)		(0.9)	(100.0)				
210～	2		4	16	22	24	4	4					76	14	90	56.85	13.94
(%)	(2.6)		(5.3)	(21.1)	(28.9)	(31.6)	(5.3)	(5.3)					(100.0)				
合計	6	4	81	519	907	960	468	190	65	9	4	8	3,221	290	3,511	61.49	13.39
(%)	(0.2)	(0.1)	(2.5)	(16.1)	(28.2)	(29.8)	(14.5)	(5.9)	(2.0)	(0.3)	(0.1)	(0.2)	(100.0)				
記載なし	32	19	91	385	684	656	343	108	39	11	6	1	2,375	3,055	5,430	59.73	14.50
(%)	(1.3)	(0.8)	(3.8)	(16.2)	(28.8)	(27.6)	(14.4)	(4.5)	(1.6)	(0.5)	(0.3)	(0.0)	(100.0)				
総計	38	23	172	904	1,591	1,616	811	298	104	20	10	9	5,596	3,345	8,941	60.74	13.90
(%)	(0.7)	(0.4)	(3.1)	(16.2)	(28.4)	(28.9)	(14.5)	(5.3)	(1.9)	(0.4)	(0.2)	(0.2)	(100.0)				
平均	194.00	147.75	129.85	133.59	129.72	127.64	120.96	126.58	120.55	141.11	96.50	137.75	128.24	131.05	128.47		
標準偏差	55.44	8.96	43.22	38.39	37.82	36.97	34.16	41.61	32.66	37.77	23.81	40.06	37.72	41.15	38.01		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査