

表 2597 体重減少量(kg) 原疾患別（施設血液透析患者 透析回数週3回 透析歴2年以上）

| 原疾患           | 0.0未満 | 0.0～   | 1.0～   | 2.0～   | 3.0～   | 4.0～  | 5.0～  | 6.0～  | 7.0～  | 8.0～  | 合計      | 記載なし  | 総計      | 平均   | 標準偏差 |
|---------------|-------|--------|--------|--------|--------|-------|-------|-------|-------|-------|---------|-------|---------|------|------|
| 慢性糸球体腎炎       | 213   | 2,619  | 14,193 | 25,239 | 12,280 | 2,684 | 517   | 108   | 20    | 40    | 57,913  | 1,534 | 59,447  | 2.45 | 1.00 |
| (%)           | (0.4) | (4.5)  | (24.5) | (43.6) | (21.2) | (4.6) | (0.9) | (0.2) | (0.0) | (0.1) | (100.0) |       |         |      |      |
| 慢性腎盂腎炎        | 15    | 102    | 513    | 748    | 312    | 69    | 16    | 2     | 2     |       | 1,779   | 38    | 1,817   | 2.32 | 0.99 |
| (%)           | (0.8) | (5.7)  | (28.8) | (42.0) | (17.5) | (3.9) | (0.9) | (0.1) | (0.1) |       | (100.0) |       |         |      |      |
| 急速進行性糸球体腎炎    | 2     | 92     | 311    | 541    | 243    | 53    | 5     | 1     |       |       | 1,248   | 33    | 1,281   | 2.33 | 0.94 |
| (%)           | (0.2) | (7.4)  | (24.9) | (43.3) | (19.5) | (4.2) | (0.4) | (0.1) |       |       | (100.0) |       |         |      |      |
| 妊娠腎/妊娠中毒症     | 1     | 35     | 283    | 483    | 150    | 18    | 4     |       | 3     |       | 977     | 22    | 999     | 2.30 | 0.82 |
| (%)           | (0.1) | (3.6)  | (29.0) | (49.4) | (15.4) | (1.8) | (0.4) |       | (0.3) |       | (100.0) |       |         |      |      |
| その他の分類不能の腎炎   | 4     | 34     | 172    | 330    | 155    | 36    | 9     | 2     |       | 1     | 743     | 20    | 763     | 2.45 | 1.01 |
| (%)           | (0.5) | (4.6)  | (23.1) | (44.4) | (20.9) | (4.8) | (1.2) | (0.3) |       | (0.1) | (100.0) |       |         |      |      |
| 多発性嚢胞腎        | 18    | 338    | 1,749  | 2,980  | 1,261  | 224   | 36    | 4     | 3     | 7     | 6,620   | 164   | 6,784   | 2.37 | 0.99 |
| (%)           | (0.3) | (5.1)  | (26.4) | (45.0) | (19.0) | (3.4) | (0.5) | (0.1) | (0.0) | (0.1) | (100.0) |       |         |      |      |
| 腎硬化症          | 71    | 820    | 3,873  | 6,189  | 2,617  | 573   | 122   | 28    | 11    | 10    | 14,314  | 363   | 14,677  | 2.36 | 0.99 |
| (%)           | (0.5) | (5.7)  | (27.1) | (43.2) | (18.3) | (4.0) | (0.9) | (0.2) | (0.1) | (0.1) | (100.0) |       |         |      |      |
| 悪性高血圧         | 8     | 53     | 293    | 539    | 351    | 122   | 42    | 7     |       | 2     | 1,417   | 48    | 1,465   | 2.69 | 1.15 |
| (%)           | (0.6) | (3.7)  | (20.7) | (38.0) | (24.8) | (8.6) | (3.0) | (0.5) |       | (0.1) | (100.0) |       |         |      |      |
| 糖尿病性腎症        | 270   | 2,899  | 13,546 | 26,352 | 16,125 | 4,597 | 984   | 191   | 49    | 84    | 65,097  | 1,753 | 66,850  | 2.60 | 1.11 |
| (%)           | (0.4) | (4.5)  | (20.8) | (40.5) | (24.8) | (7.1) | (1.5) | (0.3) | (0.1) | (0.1) | (100.0) |       |         |      |      |
| SLE腎炎         | 6     | 77     | 346    | 507    | 204    | 45    | 5     | 1     |       |       | 1,191   | 24    | 1,215   | 2.27 | 0.92 |
| (%)           | (0.5) | (6.5)  | (29.1) | (42.6) | (17.1) | (3.8) | (0.4) | (0.1) |       |       | (100.0) |       |         |      |      |
| アミロイド腎        | 4     | 15     | 74     | 98     | 25     | 7     |       |       |       |       | 223     | 7     | 230     | 2.10 | 0.88 |
| (%)           | (1.8) | (6.7)  | (33.2) | (43.9) | (11.2) | (3.1) |       |       |       |       | (100.0) |       |         |      |      |
| 痛風腎           | 4     | 34     | 127    | 314    | 155    | 32    | 15    | 4     |       | 1     | 686     | 19    | 705     | 2.57 | 1.09 |
| (%)           | (0.6) | (5.0)  | (18.5) | (45.8) | (22.6) | (4.7) | (2.2) | (0.6) |       | (0.1) | (100.0) |       |         |      |      |
| 先天性代謝異常による腎不全 | 1     | 11     | 40     | 50     | 38     | 11    |       |       |       |       | 151     | 3     | 154     | 2.40 | 1.03 |
| (%)           | (0.7) | (7.3)  | (26.5) | (33.1) | (25.2) | (7.3) |       |       |       |       | (100.0) |       |         |      |      |
| 腎・尿路結核        | 1     | 8      | 45     | 57     | 16     | 3     |       |       |       |       | 130     | 5     | 135     | 2.13 | 0.84 |
| (%)           | (0.8) | (6.2)  | (34.6) | (43.8) | (12.3) | (2.3) |       |       |       |       | (100.0) |       |         |      |      |
| 腎・尿路結石        | 2     | 29     | 93     | 152    | 59     | 9     | 2     | 1     |       |       | 347     | 7     | 354     | 2.24 | 0.94 |
| (%)           | (0.6) | (8.4)  | (26.8) | (43.8) | (17.0) | (2.6) | (0.6) | (0.3) |       |       | (100.0) |       |         |      |      |
| 腎・尿路腫瘍        | 2     | 21     | 124    | 200    | 96     | 17    | 2     |       |       |       | 462     | 11    | 473     | 2.37 | 0.88 |
| (%)           | (0.4) | (4.5)  | (26.8) | (43.3) | (20.8) | (3.7) | (0.4) |       |       |       | (100.0) |       |         |      |      |
| 閉塞性尿路障害       | 4     | 34     | 104    | 153    | 84     | 21    | 2     | 2     |       |       | 404     | 10    | 414     | 2.34 | 1.07 |
| (%)           | (1.0) | (8.4)  | (25.7) | (37.9) | (20.8) | (5.2) | (0.5) | (0.5) |       |       | (100.0) |       |         |      |      |
| 骨髄腫           | 1     | 9      | 25     | 36     | 10     | 3     |       |       |       |       | 84      | 4     | 88      | 2.11 | 0.95 |
| (%)           | (1.2) | (10.7) | (29.8) | (42.9) | (11.9) | (3.6) |       |       |       |       | (100.0) |       |         |      |      |
| 腎形成不全         | 2     | 16     | 73     | 150    | 82     | 23    | 4     | 1     |       | 1     | 352     | 7     | 359     | 2.56 | 1.13 |
| (%)           | (0.6) | (4.5)  | (20.7) | (42.6) | (23.3) | (6.5) | (1.1) | (0.3) |       | (0.3) | (100.0) |       |         |      |      |
| 不明            | 90    | 937    | 3,830  | 5,929  | 2,749  | 670   | 126   | 37    | 4     | 10    | 14,382  | 448   | 14,830  | 2.36 | 1.04 |
| (%)           | (0.6) | (6.5)  | (26.6) | (41.2) | (19.1) | (4.7) | (0.9) | (0.3) | (0.0) | (0.1) | (100.0) |       |         |      |      |
| 再導入           | 3     | 36     | 239    | 485    | 314    | 61    | 19    |       |       |       | 1,157   | 25    | 1,182   | 2.60 | 0.94 |
| (%)           | (0.3) | (3.1)  | (20.7) | (41.9) | (27.1) | (5.3) | (1.6) |       |       |       | (100.0) |       |         |      |      |
| その他           | 33    | 231    | 972    | 1,629  | 784    | 167   | 48    | 6     | 3     | 4     | 3,877   | 110   | 3,987   | 2.40 | 1.07 |
| (%)           | (0.9) | (6.0)  | (25.1) | (42.0) | (20.2) | (4.3) | (1.2) | (0.2) | (0.1) | (0.1) | (100.0) |       |         |      |      |
| 合計            | 755   | 8,450  | 41,025 | 73,161 | 38,110 | 9,445 | 1,958 | 395   | 95    | 160   | 173,554 | 4,655 | 178,209 | 2.49 | 1.05 |
| (%)           | (0.4) | (4.9)  | (23.6) | (42.2) | (22.0) | (5.4) | (1.1) | (0.2) | (0.1) | (0.1) | (100.0) |       |         |      |      |
| 記載なし          |       |        |        |        |        |       |       |       |       |       |         |       |         |      |      |
| (%)           |       |        |        |        |        |       |       |       |       |       |         |       |         |      |      |
| 総計            | 755   | 8,450  | 41,025 | 73,161 | 38,110 | 9,445 | 1,958 | 395   | 95    | 160   | 173,554 | 4,655 | 178,209 | 2.49 | 1.05 |
| (%)           | (0.4) | (4.9)  | (23.6) | (42.2) | (22.0) | (5.4) | (1.1) | (0.2) | (0.1) | (0.1) | (100.0) |       |         |      |      |

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査