

表 2598 体重減少量(kg) 医療機関別（施設血液透析患者 透析回数週3回 透析歴2年以上）

医療機関	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	2	32	83	84	30	3					234	6	240	1.92	0.94
(%)	(0.9)	(13.7)	(35.5)	(35.9)	(12.8)	(1.3)					(100.0)				
私立大学	12	74	279	392	197	45	10	2		1	1,012	7	1,019	2.32	1.06
(%)	(1.2)	(7.3)	(27.6)	(38.7)	(19.5)	(4.4)	(1.0)	(0.2)		(0.1)	(100.0)				
国立	1	20	73	143	79	8	5	1			330	1	331	2.43	0.99
(%)	(0.3)	(6.1)	(22.1)	(43.3)	(23.9)	(2.4)	(1.5)	(0.3)			(100.0)				
県市町村立	45	478	2,305	4,010	1,996	368	70	10	2	3	9,287	459	9,746	2.41	0.98
(%)	(0.5)	(5.1)	(24.8)	(43.2)	(21.5)	(4.0)	(0.8)	(0.1)	(0.0)	(0.0)	(100.0)				
社会保険	6	95	515	910	473	92	12	3		3	2,109	44	2,153	2.47	0.98
(%)	(0.3)	(4.5)	(24.4)	(43.1)	(22.4)	(4.4)	(0.6)	(0.1)		(0.1)	(100.0)				
厚生連	36	228	1,030	2,183	1,126	259	54	17	2	4	4,939	157	5,096	2.52	1.03
(%)	(0.7)	(4.6)	(20.9)	(44.2)	(22.8)	(5.2)	(1.1)	(0.3)	(0.0)	(0.1)	(100.0)				
その他公的	31	256	1,244	2,287	1,175	274	51	9	3	4	5,334	249	5,583	2.47	1.00
(%)	(0.6)	(4.8)	(23.3)	(42.9)	(22.0)	(5.1)	(1.0)	(0.2)	(0.1)	(0.1)	(100.0)				
私立総合	17	181	803	1,537	766	182	33	5	2	1	3,527	86	3,613	2.46	0.98
(%)	(0.5)	(5.1)	(22.8)	(43.6)	(21.7)	(5.2)	(0.9)	(0.1)	(0.1)	(0.0)	(100.0)				
私立	254	3,350	12,967	20,020	9,936	2,528	517	92	25	31	49,720	1,894	51,614	2.39	1.07
(%)	(0.5)	(6.7)	(26.1)	(40.3)	(20.0)	(5.1)	(1.0)	(0.2)	(0.1)	(0.1)	(100.0)				
私立診療所	351	3,736	21,726	41,595	22,332	5,686	1,206	256	61	113	97,062	1,752	98,814	2.55	1.05
(%)	(0.4)	(3.8)	(22.4)	(42.9)	(23.0)	(5.9)	(1.2)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査