

表 2606 体重減少量(kg) Body Mass Index別（施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上）

Body Mass Index	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
14未満	4	63	133	102	11	1				2	316	26	342	1.66	1.05
(%)	(1.3)	(19.9)	(42.1)	(32.3)	(3.5)	(0.3)				(0.6)	(100.0)				
14～	30	285	909	983	235	16	2	2	2	7	2,471		2,471	1.95	1.03
(%)	(1.2)	(11.5)	(36.8)	(39.8)	(9.5)	(0.6)	(0.1)	(0.1)	(0.1)	(0.3)	(100.0)				
16～	54	682	2,890	4,595	1,641	195	25	1	2	24	10,109	8	10,117	2.25	1.09
(%)	(0.5)	(6.7)	(28.6)	(45.5)	(16.2)	(1.9)	(0.2)	(0.0)	(0.0)	(0.2)	(100.0)				
18～	101	1,024	4,832	9,343	4,689	715	108	13	6	24	20,855	9	20,864	2.43	1.00
(%)	(0.5)	(4.9)	(23.2)	(44.8)	(22.5)	(3.4)	(0.5)	(0.1)	(0.0)	(0.1)	(100.0)				
20～	113	956	4,697	11,146	6,871	1,360	210	22	7	18	25,400	8	25,408	2.59	0.97
(%)	(0.4)	(3.8)	(18.5)	(43.9)	(27.1)	(5.4)	(0.8)	(0.1)	(0.0)	(0.1)	(100.0)				
22～	65	638	3,101	8,278	6,227	1,592	229	41	8	6	20,185	5	20,190	2.74	0.98
(%)	(0.3)	(3.2)	(15.4)	(41.0)	(30.8)	(7.9)	(1.1)	(0.2)	(0.0)	(0.0)	(100.0)				
24～	35	264	1,475	4,297	4,191	1,381	269	33	8	9	11,962	2	11,964	2.94	1.04
(%)	(0.3)	(2.2)	(12.3)	(35.9)	(35.0)	(11.5)	(2.2)	(0.3)	(0.1)	(0.1)	(100.0)				
26～	9	135	575	1,841	2,334	1,001	262	47	5	2	6,211	3	6,214	3.18	1.07
(%)	(0.1)	(2.2)	(9.3)	(29.6)	(37.6)	(16.1)	(4.2)	(0.8)	(0.1)	(0.0)	(100.0)				
28～	9	52	198	742	1,095	721	179	44	5	2	3,047	3	3,050	3.43	1.12
(%)	(0.3)	(1.7)	(6.5)	(24.4)	(35.9)	(23.7)	(5.9)	(1.4)	(0.2)	(0.1)	(100.0)				
30～	4	23	110	330	770	737	304	91	17	8	2,394	4	2,398	3.91	1.26
(%)	(0.2)	(1.0)	(4.6)	(13.8)	(32.2)	(30.8)	(12.7)	(3.8)	(0.7)	(0.3)	(100.0)				
合計	424	4,122	18,920	41,657	28,064	7,719	1,588	294	60	102	102,950	68	103,018	2.67	1.08
(%)	(0.4)	(4.0)	(18.4)	(40.5)	(27.3)	(7.5)	(1.5)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	33	378	1,325	2,377	1,603	600	221	82	23	14	6,656	2,841	9,497	2.73	1.29
(%)	(0.5)	(5.7)	(19.9)	(35.7)	(24.1)	(9.0)	(3.3)	(1.2)	(0.3)	(0.2)	(100.0)				
総計	457	4,500	20,245	44,034	29,667	8,319	1,809	376	83	116	109,606	2,909	112,515	2.67	1.09
(%)	(0.4)	(4.1)	(18.5)	(40.2)	(27.1)	(7.6)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	20.60	20.21	20.49	21.26	22.53	24.40	25.84	27.20	25.82	20.57	21.74	17.05	21.74		
標準偏差	3.30	3.24	3.07	3.06	3.36	3.88	4.26	4.37	5.44	4.61	3.48	7.75	3.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査