

表 2609 体重減少量(kg) 透析前クレアチニン濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前クレアチニン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
2未満	3	18	35	12	7		1				76		76	1.54	1.07
(%)	(3.9)	(23.7)	(46.1)	(15.8)	(9.2)		(1.3)				(100.0)				
2～	42	421	723	376	56	9	4	1	1	2	1,635	21	1,656	1.49	1.13
(%)	(2.6)	(25.7)	(44.2)	(23.0)	(3.4)	(0.6)	(0.2)	(0.1)	(0.1)	(0.1)	(100.0)				
4～	137	1,391	3,237	2,371	486	64	13	3	1	8	7,711	84	7,795	1.71	0.98
(%)	(1.8)	(18.0)	(42.0)	(30.7)	(6.3)	(0.8)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
6～	194	2,151	8,390	9,673	2,828	465	88	14	7	21	23,831	175	24,006	2.07	0.96
(%)	(0.8)	(9.0)	(35.2)	(40.6)	(11.9)	(2.0)	(0.4)	(0.1)	(0.0)	(0.1)	(100.0)				
8～	158	2,195	12,658	20,931	8,172	1,418	257	41	19	44	45,893	222	46,115	2.34	0.95
(%)	(0.3)	(4.8)	(27.6)	(45.6)	(17.8)	(3.1)	(0.6)	(0.1)	(0.0)	(0.1)	(100.0)				
10～	106	1,349	9,942	21,840	11,422	2,529	439	85	16	45	47,773	261	48,034	2.57	0.98
(%)	(0.2)	(2.8)	(20.8)	(45.7)	(23.9)	(5.3)	(0.9)	(0.2)	(0.0)	(0.1)	(100.0)				
12～	63	579	4,236	12,288	9,126	2,594	535	114	24	20	29,579	175	29,754	2.83	1.00
(%)	(0.2)	(2.0)	(14.3)	(41.5)	(30.9)	(8.8)	(1.8)	(0.4)	(0.1)	(0.1)	(100.0)				
14～	34	183	1,214	4,187	4,280	1,584	384	76	19	13	11,974	62	12,036	3.07	1.07
(%)	(0.3)	(1.5)	(10.1)	(35.0)	(35.7)	(13.2)	(3.2)	(0.6)	(0.2)	(0.1)	(100.0)				
16～	9	69	299	1,101	1,552	736	230	58	6	7	4,067	33	4,100	3.33	1.18
(%)	(0.2)	(1.7)	(7.4)	(27.1)	(38.2)	(18.1)	(5.7)	(1.4)	(0.1)	(0.2)	(100.0)				
合計	746	8,356	40,734	72,779	37,929	9,399	1,951	392	93	160	172,539	1,033	173,572	2.49	1.05
(%)	(0.4)	(4.8)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	9	94	291	382	181	46	7	3	2		1,015	3,622	4,637	2.26	1.08
(%)	(0.9)	(9.3)	(28.7)	(37.6)	(17.8)	(4.5)	(0.7)	(0.3)	(0.2)		(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	8.38	8.27	9.25	10.27	11.33	12.15	12.63	12.82	11.73	10.44	10.29	10.00	10.29		
標準偏差	3.20	2.91	2.59	2.53	2.63	2.69	3.05	2.91	2.90	3.00	2.78	3.19	2.79		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査