

表 2614 体重減少量(kg) equilibrated Kt/V(eKt/V)別(施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上)

equilibrated Kt/V (eKt/V)	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.4未満	5	9	33	32	18	5	1				103		103	2.13	1.21
(%)	(4.9)	(8.7)	(32.0)	(31.1)	(17.5)	(4.9)	(1.0)				(100.0)				
0.4～	6	62	69	70	43	6	2			1	259		259	1.89	1.52
(%)	(2.3)	(23.9)	(26.6)	(27.0)	(16.6)	(2.3)	(0.8)			(0.4)	(100.0)				
0.6～	47	332	673	735	293	108	40	12	1		2,241		2,241	2.08	1.23
(%)	(2.1)	(14.8)	(30.0)	(32.8)	(13.1)	(4.8)	(1.8)	(0.5)	(0.0)		(100.0)				
0.8～	110	925	2,816	4,399	2,592	896	241	62	9	4	12,054		12,054	2.48	1.17
(%)	(0.9)	(7.7)	(23.4)	(36.5)	(21.5)	(7.4)	(2.0)	(0.5)	(0.1)	(0.0)	(100.0)				
1.0～	135	1,309	5,908	12,999	9,510	2,717	520	98	22	7	33,225		33,225	2.69	1.03
(%)	(0.4)	(3.9)	(17.8)	(39.1)	(28.6)	(8.2)	(1.6)	(0.3)	(0.1)	(0.0)	(100.0)				
1.2～	84	944	5,855	14,921	10,062	2,509	524	98	20	17	35,034		35,034	2.72	0.97
(%)	(0.2)	(2.7)	(16.7)	(42.6)	(28.7)	(7.2)	(1.5)	(0.3)	(0.1)	(0.0)	(100.0)				
1.4～	26	370	2,553	6,288	4,067	1,078	227	46	14	19	14,688	1	14,689	2.72	1.02
(%)	(0.2)	(2.5)	(17.4)	(42.8)	(27.7)	(7.3)	(1.5)	(0.3)	(0.1)	(0.1)	(100.0)				
1.6～	3	96	675	1,715	1,166	360	99	29	7	25	4,175		4,175	2.86	1.30
(%)	(0.1)	(2.3)	(16.2)	(41.1)	(27.9)	(8.6)	(2.4)	(0.7)	(0.2)	(0.6)	(100.0)				
1.8～	2	38	196	408	317	122	33	10	4	13	1,143	1	1,144	2.94	1.56
(%)	(0.2)	(3.3)	(17.1)	(35.7)	(27.7)	(10.7)	(2.9)	(0.9)	(0.3)	(1.1)	(100.0)				
2.0～	3	14	77	228	169	83	28	5	1	22	630	17	647	3.45	2.98
(%)	(0.5)	(2.2)	(12.2)	(36.2)	(26.8)	(13.2)	(4.4)	(0.8)	(0.2)	(3.5)	(100.0)				
合計	421	4,099	18,855	41,795	28,237	7,884	1,715	360	78	108	103,552	19	103,571	2.68	1.09
(%)	(0.4)	(4.0)	(18.2)	(40.4)	(27.3)	(7.6)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
記載なし	36	401	1,390	2,239	1,430	435	94	16	5	8	6,054	2,890	8,944	2.53	1.21
(%)	(0.6)	(6.6)	(23.0)	(37.0)	(23.6)	(7.2)	(1.6)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	457	4,500	20,245	44,034	29,667	8,319	1,809	376	83	116	109,606	2,909	112,515	2.67	1.09
(%)	(0.4)	(4.1)	(18.5)	(40.2)	(27.1)	(7.6)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
平均	1.06	1.12	1.20	1.24	1.24	1.24	1.24	1.24	1.31	1.66	1.22	2.58	1.23		
標準偏差	0.26	0.26	0.24	0.23	0.22	0.25	0.27	0.28	0.28	0.42	0.24	0.39	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査