

表 2615 体重減少量(kg) equilibrated Kt/V(eKt/V)別(施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上)

equilibrated Kt/V (eKt/V)	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.4未満		7	10	10	3	1					31		31	1.85	0.98
(%)		(22.6)	(32.3)	(32.3)	(9.7)	(3.2)					(100.0)				
0.4～	6	28	36	34	3						107		107	1.43	0.92
(%)	(5.6)	(26.2)	(33.6)	(31.8)	(2.8)						(100.0)				
0.6～	19	115	248	175	29	10	1				597		597	1.63	0.93
(%)	(3.2)	(19.3)	(41.5)	(29.3)	(4.9)	(1.7)	(0.2)				(100.0)				
0.8～	38	426	1,195	983	284	54	6				2,986		2,986	1.86	0.94
(%)	(1.3)	(14.3)	(40.0)	(32.9)	(9.5)	(1.8)	(0.2)				(100.0)				
1.0～	78	816	3,243	3,685	1,107	215	16	6	1	3	9,170		9,170	2.07	0.91
(%)	(0.9)	(8.9)	(35.4)	(40.2)	(12.1)	(2.3)	(0.2)	(0.1)	(0.0)	(0.0)	(100.0)				
1.2～	64	919	5,201	7,705	2,477	319	43	3	1	2	16,734		16,734	2.21	0.83
(%)	(0.4)	(5.5)	(31.1)	(46.0)	(14.8)	(1.9)	(0.3)	(0.0)	(0.0)	(0.0)	(100.0)				
1.4～	36	732	4,963	8,270	2,334	257	33	4	2	3	16,634		16,634	2.24	0.78
(%)	(0.2)	(4.4)	(29.8)	(49.7)	(14.0)	(1.5)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
1.6～	24	354	2,975	4,399	1,121	115	19	3	4	8	9,022		9,022	2.20	0.81
(%)	(0.3)	(3.9)	(33.0)	(48.8)	(12.4)	(1.3)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
1.8～	6	120	1,045	1,594	398	49	12		3	5	3,232	1	3,233	2.22	0.85
(%)	(0.2)	(3.7)	(32.3)	(49.3)	(12.3)	(1.5)	(0.4)		(0.1)	(0.2)	(100.0)				
2.0～	4	70	578	825	298	41	7	1	1	14	1,839	2	1,841	2.38	1.50
(%)	(0.2)	(3.8)	(31.4)	(44.9)	(16.2)	(2.2)	(0.4)	(0.1)	(0.1)	(0.8)	(100.0)				
合計	275	3,587	19,494	27,680	8,054	1,061	137	17	12	35	60,352	3	60,355	2.18	0.87
(%)	(0.5)	(5.9)	(32.3)	(45.9)	(13.3)	(1.8)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
記載なし	23	363	1,286	1,447	389	65	12	2		9	3,596	1,743	5,339	2.07	1.19
(%)	(0.6)	(10.1)	(35.8)	(40.2)	(10.8)	(1.8)	(0.3)	(0.1)		(0.3)	(100.0)				
総計	298	3,950	20,780	29,127	8,443	1,126	149	19	12	44	63,948	1,746	65,694	2.17	0.89
(%)	(0.5)	(6.2)	(32.5)	(45.5)	(13.2)	(1.8)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
平均	1.22	1.29	1.40	1.43	1.43	1.38	1.46	1.44	1.67	1.87	1.41	2.42	1.41		
標準偏差	0.32	0.31	0.30	0.28	0.28	0.30	0.31	0.37	0.30	0.44	0.29	0.54	0.29		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査