

表 2616 体重減少量(kg) nPCR(g/kg/day)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

nPCR	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.4未満	24	117	117	79	26	5		1			369		369	1.42	1.13
(%)	(6.5)	(31.7)	(31.7)	(21.4)	(7.0)	(1.4)		(0.3)			(100.0)				
0.4～	214	1,645	3,360	2,453	634	95	11	4	2		8,418		8,418	1.68	0.96
(%)	(2.5)	(19.5)	(39.9)	(29.1)	(7.5)	(1.1)	(0.1)	(0.0)	(0.0)		(100.0)				
0.6～	320	3,511	15,401	21,010	8,642	1,720	275	54	13	9	50,955	1	50,956	2.25	0.96
(%)	(0.6)	(6.9)	(30.2)	(41.2)	(17.0)	(3.4)	(0.5)	(0.1)	(0.0)	(0.0)	(100.0)				
0.8～	117	1,945	14,582	31,508	17,126	4,168	847	139	39	36	70,507	4	70,511	2.60	0.95
(%)	(0.2)	(2.8)	(20.7)	(44.7)	(24.3)	(5.9)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
1.0～	17	405	4,231	12,281	8,179	2,377	547	123	23	62	28,245	3	28,248	2.83	1.06
(%)	(0.1)	(1.4)	(15.0)	(43.5)	(29.0)	(8.4)	(1.9)	(0.4)	(0.1)	(0.2)	(100.0)				
1.2～	3	58	602	1,996	1,530	505	136	51	9	23	4,913	3	4,916	3.01	1.27
(%)	(0.1)	(1.2)	(12.3)	(40.6)	(31.1)	(10.3)	(2.8)	(1.0)	(0.2)	(0.5)	(100.0)				
1.4～	1	9	66	181	162	71	36	5	4	9	544	11	555	3.34	2.21
(%)	(0.2)	(1.7)	(12.1)	(33.3)	(29.8)	(13.1)	(6.6)	(0.9)	(0.7)	(1.7)	(100.0)				
1.6～		1	15	18	24	12				5	75	4	79	3.80	3.80
(%)		(1.3)	(20.0)	(24.0)	(32.0)	(16.0)				(6.7)	(100.0)				
1.8～			2	7						3	12	13	25	6.00	7.19
(%)			(16.7)	(58.3)						(25.0)	(100.0)				
2.0～			3	1	2	1	1			2	10	28	38	6.74	8.25
(%)			(30.0)	(10.0)	(20.0)	(10.0)	(10.0)			(20.0)	(100.0)				
合計	696	7,691	38,379	69,534	36,325	8,954	1,853	377	90	149	164,048	67	164,115	2.49	1.04
(%)	(0.4)	(4.7)	(23.4)	(42.4)	(22.1)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	59	759	2,646	3,627	1,785	491	105	18	5	11	9,506	4,588	14,094	2.35	1.17
(%)	(0.6)	(8.0)	(27.8)	(38.2)	(18.8)	(5.2)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	0.67	0.72	0.81	0.87	0.90	0.93	0.96	0.99	0.98	1.14	0.86	1.89	0.86		
標準偏差	0.16	0.17	0.17	0.16	0.16	0.17	0.18	0.19	0.21	0.28	0.17	0.53	0.18		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査