

表 2619 体重減少量(kg) %クレアチニン産生速度(%)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	266	2,286	4,904	3,744	1,075	210	52	9	2	2	12,550		12,550	1.76	1.00
(%)	(2.1)	(18.2)	(39.1)	(29.8)	(8.6)	(1.7)	(0.4)	(0.1)	(0.0)	(0.0)	(100.0)				
60～	92	953	3,007	3,361	1,286	334	69	20	1	2	9,125		9,125	2.12	1.03
(%)	(1.0)	(10.4)	(33.0)	(36.8)	(14.1)	(3.7)	(0.8)	(0.2)	(0.0)	(0.0)	(100.0)				
70～	94	986	4,100	5,583	2,442	605	120	16	6	1	13,953	1	13,954	2.29	1.00
(%)	(0.7)	(7.1)	(29.4)	(40.0)	(17.5)	(4.3)	(0.9)	(0.1)	(0.0)	(0.0)	(100.0)				
80～	79	951	5,036	8,365	4,031	932	214	49	14	12	19,683		19,683	2.43	1.00
(%)	(0.4)	(4.8)	(25.6)	(42.5)	(20.5)	(4.7)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
90～	59	865	5,749	10,801	5,566	1,445	299	47	16	11	24,858		24,858	2.53	0.98
(%)	(0.2)	(3.5)	(23.1)	(43.5)	(22.4)	(5.8)	(1.2)	(0.2)	(0.1)	(0.0)	(100.0)				
100～	40	703	5,523	11,661	6,593	1,657	320	77	11	10	26,595		26,595	2.61	0.95
(%)	(0.2)	(2.6)	(20.8)	(43.8)	(24.8)	(6.2)	(1.2)	(0.3)	(0.0)	(0.0)	(100.0)				
110～	33	467	4,461	10,812	6,131	1,525	319	73	20	12	23,853	1	23,854	2.67	0.95
(%)	(0.1)	(2.0)	(18.7)	(45.3)	(25.7)	(6.4)	(1.3)	(0.3)	(0.1)	(0.1)	(100.0)				
120～	17	259	3,063	7,818	4,717	1,166	233	45	8	12	17,338		17,338	2.71	0.94
(%)	(0.1)	(1.5)	(17.7)	(45.1)	(27.2)	(6.7)	(1.3)	(0.3)	(0.0)	(0.1)	(100.0)				
130～	6	121	1,516	4,421	2,708	666	130	23	4	13	9,608		9,608	2.75	0.97
(%)	(0.1)	(1.3)	(15.8)	(46.0)	(28.2)	(6.9)	(1.4)	(0.2)	(0.0)	(0.1)	(100.0)				
140～	10	89	1,006	2,949	1,765	412	94	18	8	74	6,425	24	6,449	2.87	1.61
(%)	(0.2)	(1.4)	(15.7)	(45.9)	(27.5)	(6.4)	(1.5)	(0.3)	(0.1)	(1.2)	(100.0)				
合計	696	7,680	38,365	69,515	36,314	8,952	1,850	377	90	149	163,988	26	164,014	2.49	1.04
(%)	(0.4)	(4.7)	(23.4)	(42.4)	(22.1)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	59	770	2,660	3,646	1,796	493	108	18	5	11	9,566	4,629	14,195	2.34	1.17
(%)	(0.6)	(8.0)	(27.8)	(38.1)	(18.8)	(5.2)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	69.91	75.98	91.78	101.09	104.78	105.03	104.62	104.31	106.29	142.89	98.72	219.40	98.74		
標準偏差	29.59	28.69	27.30	24.42	22.54	21.99	22.93	21.87	23.93	43.19	25.91	50.77	25.96		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査