

表 2621 体重減少量(kg) %クレアチニン産生速度(%)別 (施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	113	1,024	2,296	1,399	264	25	4		1		5,126		5,126	1.60	0.86
(%)	(2.2)	(20.0)	(44.8)	(27.3)	(5.2)	(0.5)	(0.1)		(0.0)		(100.0)				
60～	33	409	1,379	1,188	264	55	8	2		1	3,339		3,339	1.88	0.89
(%)	(1.0)	(12.2)	(41.3)	(35.6)	(7.9)	(1.6)	(0.2)	(0.1)		(0.0)	(100.0)				
70～	37	462	1,923	1,984	544	80	12	2	1		5,045		5,045	2.01	0.87
(%)	(0.7)	(9.2)	(38.1)	(39.3)	(10.8)	(1.6)	(0.2)	(0.0)	(0.0)		(100.0)				
80～	31	444	2,456	3,151	899	115	20	3		5	7,124		7,124	2.14	0.85
(%)	(0.4)	(6.2)	(34.5)	(44.2)	(12.6)	(1.6)	(0.3)	(0.0)		(0.1)	(100.0)				
90～	26	418	2,859	4,126	1,233	186	18	2		1	8,869		8,869	2.21	0.81
(%)	(0.3)	(4.7)	(32.2)	(46.5)	(13.9)	(2.1)	(0.2)	(0.0)		(0.0)	(100.0)				
100～	18	369	2,948	4,693	1,510	210	23	4	1	1	9,777		9,777	2.28	0.80
(%)	(0.2)	(3.8)	(30.2)	(48.0)	(15.4)	(2.1)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
110～	6	219	2,475	4,583	1,332	146	20	1	4	4	8,790	1	8,791	2.32	0.75
(%)	(0.1)	(2.5)	(28.2)	(52.1)	(15.2)	(1.7)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
120～	7	139	1,737	3,403	1,087	125	14	2	3	2	6,519		6,519	2.36	0.76
(%)	(0.1)	(2.1)	(26.6)	(52.2)	(16.7)	(1.9)	(0.2)	(0.0)	(0.0)	(0.0)	(100.0)				
130～	2	63	860	1,922	566	80	12	1		3	3,509		3,509	2.40	0.83
(%)	(0.1)	(1.8)	(24.5)	(54.8)	(16.1)	(2.3)	(0.3)	(0.0)		(0.1)	(100.0)				
140～	2	35	572	1,249	361	42	6		2	22	2,291	7	2,298	2.49	1.42
(%)	(0.1)	(1.5)	(25.0)	(54.5)	(15.8)	(1.8)	(0.3)		(0.1)	(1.0)	(100.0)				
合計	275	3,582	19,505	27,698	8,060	1,064	137	17	12	39	60,389	8	60,397	2.18	0.88
(%)	(0.5)	(5.9)	(32.3)	(45.9)	(13.3)	(1.8)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
記載なし	23	368	1,275	1,429	383	62	12	2		5	3,559	1,738	5,297	2.04	1.11
(%)	(0.6)	(10.3)	(35.8)	(40.2)	(10.8)	(1.7)	(0.3)	(0.1)		(0.1)	(100.0)				
総計	298	3,950	20,780	29,127	8,443	1,126	149	19	12	44	63,948	1,746	65,694	2.17	0.89
(%)	(0.5)	(6.2)	(32.5)	(45.5)	(13.2)	(1.8)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
平均	67.50	77.18	93.60	102.54	104.43	103.27	102.61	95.61	116.25	151.12	98.29	200.14	98.30		
標準偏差	28.26	28.47	27.48	24.29	22.56	22.49	27.35	19.17	32.47	48.78	26.46	46.61	26.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査