

表 2622 体重減少量(kg) 透析前アルブミン濃度(g/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前アルブミン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
1.5未満	5	21	31	8	3		2				70	3	73	1.29	1.10
(%)	(7.1)	(30.0)	(44.3)	(11.4)	(4.3)		(2.9)				(100.0)				
1.5～	14	138	185	96	24	6					463	11	474	1.41	0.93
(%)	(3.0)	(29.8)	(40.0)	(20.7)	(5.2)	(1.3)					(100.0)				
2.0～	48	483	856	562	93	13	3			1	2,059	35	2,094	1.56	0.91
(%)	(2.3)	(23.5)	(41.6)	(27.3)	(4.5)	(0.6)	(0.1)			(0.0)	(100.0)				
2.5～	91	1,098	3,301	3,103	831	137	19	8	3	8	8,599	97	8,696	1.93	1.00
(%)	(1.1)	(12.8)	(38.4)	(36.1)	(9.7)	(1.6)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
3.0～	157	2,224	11,046	17,302	7,238	1,421	251	50	19	32	39,740	266	40,006	2.34	0.99
(%)	(0.4)	(5.6)	(27.8)	(43.5)	(18.2)	(3.6)	(0.6)	(0.1)	(0.0)	(0.1)	(100.0)				
3.5～	248	2,950	18,030	36,783	20,115	4,914	1,052	199	50	75	84,416	478	84,894	2.57	1.02
(%)	(0.3)	(3.5)	(21.4)	(43.6)	(23.8)	(5.8)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
4.0～	151	1,176	6,088	12,866	8,352	2,554	540	116	20	35	31,898	220	32,118	2.67	1.09
(%)	(0.5)	(3.7)	(19.1)	(40.3)	(26.2)	(8.0)	(1.7)	(0.4)	(0.1)	(0.1)	(100.0)				
4.5～	21	100	345	678	510	138	32	10		3	1,837	10	1,847	2.61	1.14
(%)	(1.1)	(5.4)	(18.8)	(36.9)	(27.8)	(7.5)	(1.7)	(0.5)		(0.2)	(100.0)				
5.0～	3	8	20	49	28	9	4	1			122		122	2.59	1.19
(%)	(2.5)	(6.6)	(16.4)	(40.2)	(23.0)	(7.4)	(3.3)	(0.8)			(100.0)				
合計	738	8,198	39,902	71,447	37,194	9,192	1,903	384	92	154	169,204	1,120	170,324	2.49	1.05
(%)	(0.4)	(4.8)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	17	252	1,123	1,714	916	253	55	11	3	6	4,350	3,535	7,885	2.45	1.16
(%)	(0.4)	(5.8)	(25.8)	(39.4)	(21.1)	(5.8)	(1.3)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	3.45	3.39	3.52	3.62	3.70	3.76	3.78	3.79	3.66	3.65	3.61	3.53	3.61		
標準偏差	0.67	0.59	0.47	0.40	0.36	0.35	0.35	0.36	0.34	0.41	0.43	0.52	0.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査