

表 2623 体重減少量(kg) 透析前CRP濃度(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前CRP濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.2未満	280	2,996	18,038	35,577	18,749	4,410	797	138	28	60	81,073	433	81,506	2.53	1.00
(%)	(0.3)	(3.7)	(22.2)	(43.9)	(23.1)	(5.4)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
0.2～	104	1,224	6,449	11,522	6,263	1,652	354	72	19	31	27,690	150	27,840	2.53	1.05
(%)	(0.4)	(4.4)	(23.3)	(41.6)	(22.6)	(6.0)	(1.3)	(0.3)	(0.1)	(0.1)	(100.0)				
0.5～	54	798	3,485	5,765	2,936	840	191	39	9	14	14,131	115	14,246	2.47	1.09
(%)	(0.4)	(5.6)	(24.7)	(40.8)	(20.8)	(5.9)	(1.4)	(0.3)	(0.1)	(0.1)	(100.0)				
1.0～	61	719	2,516	3,533	1,714	453	109	29	17	7	9,158	87	9,245	2.35	1.10
(%)	(0.7)	(7.9)	(27.5)	(38.6)	(18.7)	(4.9)	(1.2)	(0.3)	(0.2)	(0.1)	(100.0)				
2.0～	48	553	1,678	2,105	897	212	64	22	7	5	5,591	52	5,643	2.24	1.14
(%)	(0.9)	(9.9)	(30.0)	(37.6)	(16.0)	(3.8)	(1.1)	(0.4)	(0.1)	(0.1)	(100.0)				
4.0～	31	230	639	665	295	67	16	2	2	2	1,949	27	1,976	2.12	1.18
(%)	(1.6)	(11.8)	(32.8)	(34.1)	(15.1)	(3.4)	(0.8)	(0.1)	(0.1)	(0.1)	(100.0)				
6.0～	15	153	312	302	116	23	11	2	1	3	938	11	949	1.99	1.18
(%)	(1.6)	(16.3)	(33.3)	(32.2)	(12.4)	(2.5)	(1.2)	(0.2)	(0.1)	(0.3)	(100.0)				
8.0～	14	90	164	167	62	14	6	4			521	11	532	1.94	1.18
(%)	(2.7)	(17.3)	(31.5)	(32.1)	(11.9)	(2.7)	(1.2)	(0.8)			(100.0)				
10.0～	17	108	209	205	82	19	5	1			646	5	651	1.91	1.11
(%)	(2.6)	(16.7)	(32.4)	(31.7)	(12.7)	(2.9)	(0.8)	(0.2)			(100.0)				
15.0～	15	82	131	134	59	16	3				440	9	449	1.93	1.15
(%)	(3.4)	(18.6)	(29.8)	(30.5)	(13.4)	(3.6)	(0.7)				(100.0)				
合計	639	6,953	33,621	59,975	31,173	7,706	1,556	309	83	122	142,137	900	143,037	2.48	1.05
(%)	(0.4)	(4.9)	(23.7)	(42.2)	(21.9)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	116	1,497	7,404	13,186	6,937	1,739	402	86	12	38	31,417	3,755	35,172	2.50	1.08
(%)	(0.4)	(4.8)	(23.6)	(42.0)	(22.1)	(5.5)	(1.3)	(0.3)	(0.0)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	1.89	1.41	0.78	0.56	0.49	0.50	0.61	0.76	0.84	0.62	0.64	1.16	0.64		
標準偏差	4.00	3.20	2.16	1.81	1.58	1.57	1.41	1.55	1.10	1.23	1.95	3.46	1.97		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査