

表 2625 体重減少量(kg) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
3未満	97	1,055	2,435	2,366	705	136	27	1	1	4	6,827	62	6,889	1.90	1.05
(%)	(1.4)	(15.5)	(35.7)	(34.7)	(10.3)	(2.0)	(0.4)	(0.0)	(0.0)	(0.1)	(100.0)				
3.0～	64	809	2,719	3,180	1,145	209	29	9	4	10	8,178	58	8,236	2.11	1.04
(%)	(0.8)	(9.9)	(33.2)	(38.9)	(14.0)	(2.6)	(0.4)	(0.1)	(0.0)	(0.1)	(100.0)				
3.5～	95	1,076	4,321	5,860	2,224	419	66	17	5	11	14,094	100	14,194	2.21	0.99
(%)	(0.7)	(7.6)	(30.7)	(41.6)	(15.8)	(3.0)	(0.5)	(0.1)	(0.0)	(0.1)	(100.0)				
4.0～	112	1,263	6,075	9,172	3,819	717	118	28	6	17	21,327	125	21,452	2.31	0.99
(%)	(0.5)	(5.9)	(28.5)	(43.0)	(17.9)	(3.4)	(0.6)	(0.1)	(0.0)	(0.1)	(100.0)				
4.5～	120	1,171	6,551	11,478	5,292	1,065	171	28	10	19	25,905	173	26,078	2.41	0.97
(%)	(0.5)	(4.5)	(25.3)	(44.3)	(20.4)	(4.1)	(0.7)	(0.1)	(0.0)	(0.1)	(100.0)				
5.0～	89	1,032	6,189	11,839	5,804	1,199	210	38	5	27	26,432	192	26,624	2.48	0.99
(%)	(0.3)	(3.9)	(23.4)	(44.8)	(22.0)	(4.5)	(0.8)	(0.1)	(0.0)	(0.1)	(100.0)				
5.5～	44	731	4,718	10,326	5,535	1,313	248	38	13	22	22,988	134	23,122	2.58	1.01
(%)	(0.2)	(3.2)	(20.5)	(44.9)	(24.1)	(5.7)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
6.0～	10	112	780	1,705	997	281	43	11	2	4	3,945	21	3,966	2.64	1.00
(%)	(0.3)	(2.8)	(19.8)	(43.2)	(25.3)	(7.1)	(1.1)	(0.3)	(0.1)	(0.1)	(100.0)				
6.1～	24	370	2,483	5,640	3,434	901	199	33	5	13	13,102	63	13,165	2.67	1.04
(%)	(0.2)	(2.8)	(19.0)	(43.0)	(26.2)	(6.9)	(1.5)	(0.3)	(0.0)	(0.1)	(100.0)				
6.5～	20	270	1,884	4,665	3,261	953	190	39	7	12	11,301	61	11,362	2.76	1.02
(%)	(0.2)	(2.4)	(16.7)	(41.3)	(28.9)	(8.4)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
7.0～	14	168	1,049	2,740	2,151	693	171	30	7	7	7,030	50	7,080	2.85	1.05
(%)	(0.2)	(2.4)	(14.9)	(39.0)	(30.6)	(9.9)	(2.4)	(0.4)	(0.1)	(0.1)	(100.0)				
7.5～	19	90	548	1,549	1,316	495	115	18	10	4	4,164	34	4,198	2.94	1.07
(%)	(0.5)	(2.2)	(13.2)	(37.2)	(31.6)	(11.9)	(2.8)	(0.4)	(0.2)	(0.1)	(100.0)				
8.0～	7	54	288	788	880	306	101	35	9	3	2,471	14	2,485	3.08	1.15
(%)	(0.3)	(2.2)	(11.7)	(31.9)	(35.6)	(12.4)	(4.1)	(1.4)	(0.4)	(0.1)	(100.0)				
8.5～	4	38	209	516	485	238	78	20	4	1	1,593	8	1,601	3.08	1.21
(%)	(0.3)	(2.4)	(13.1)	(32.4)	(30.4)	(14.9)	(4.9)	(1.3)	(0.3)	(0.1)	(100.0)				
9.0～	18	58	294	728	752	435	175	45	5	4	2,514	23	2,537	3.21	1.31
(%)	(0.7)	(2.3)	(11.7)	(29.0)	(29.9)	(17.3)	(7.0)	(1.8)	(0.2)	(0.2)	(100.0)				
合計	737	8,297	40,543	72,552	37,800	9,360	1,941	390	93	158	171,871	1,118	172,989	2.49	1.05
(%)	(0.4)	(4.8)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	18	153	482	609	310	85	17	5	2	2	1,683	3,537	5,220	2.29	1.17
(%)	(1.1)	(9.1)	(28.6)	(36.2)	(18.4)	(5.1)	(1.0)	(0.3)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	4.60	4.50	4.87	5.20	5.56	5.96	6.37	6.63	6.28	5.40	5.22	5.15	5.22		
標準偏差	1.64	1.46	1.33	1.32	1.42	1.61	1.81	2.03	1.86	1.55	1.42	1.54	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査