

表 2628 体重減少量(kg) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前ヘモグロビン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
5未満	3	9	52	91	39	7	3	1			205	5	210	2.40	1.01
(%)	(1.5)	(4.4)	(25.4)	(44.4)	(19.0)	(3.4)	(1.5)	(0.5)			(100.0)				
5～	2	23	52	107	54	5		1			244	7	251	2.31	0.97
(%)	(0.8)	(9.4)	(21.3)	(43.9)	(22.1)	(2.0)		(0.4)			(100.0)				
6～	9	64	171	233	95	27	5	3			607	10	617	2.20	1.07
(%)	(1.5)	(10.5)	(28.2)	(38.4)	(15.7)	(4.4)	(0.8)	(0.5)			(100.0)				
7～	21	253	706	879	419	100	22	8	1	2	2,411	34	2,445	2.23	1.10
(%)	(0.9)	(10.5)	(29.3)	(36.5)	(17.4)	(4.1)	(0.9)	(0.3)	(0.0)	(0.1)	(100.0)				
8～	50	631	2,313	3,369	1,521	409	77	22	10	5	8,407	90	8,497	2.33	1.05
(%)	(0.6)	(7.5)	(27.5)	(40.1)	(18.1)	(4.9)	(0.9)	(0.3)	(0.1)	(0.1)	(100.0)				
9～	119	1,534	7,285	11,943	5,744	1,404	293	58	13	30	28,423	185	28,608	2.42	1.05
(%)	(0.4)	(5.4)	(25.6)	(42.0)	(20.2)	(4.9)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
10～	231	2,562	13,914	25,550	13,253	3,108	621	130	32	56	59,457	312	59,769	2.50	1.03
(%)	(0.4)	(4.3)	(23.4)	(43.0)	(22.3)	(5.2)	(1.0)	(0.2)	(0.1)	(0.1)	(100.0)				
11～	194	2,179	10,940	20,448	10,926	2,721	571	98	22	36	48,135	280	48,415	2.51	1.04
(%)	(0.4)	(4.5)	(22.7)	(42.5)	(22.7)	(5.7)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
12～	78	764	3,725	7,187	4,035	1,038	213	42	10	18	17,110	98	17,208	2.54	1.07
(%)	(0.5)	(4.5)	(21.8)	(42.0)	(23.6)	(6.1)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
13～	28	188	829	1,702	1,035	326	86	17	2	7	4,220	40	4,260	2.64	1.26
(%)	(0.7)	(4.5)	(19.6)	(40.3)	(24.5)	(7.7)	(2.0)	(0.4)	(0.0)	(0.2)	(100.0)				
14～	9	76	302	562	399	136	43	8	2	1	1,538	13	1,551	2.69	1.17
(%)	(0.6)	(4.9)	(19.6)	(36.5)	(25.9)	(8.8)	(2.8)	(0.5)	(0.1)	(0.1)	(100.0)				
合計	744	8,283	40,289	72,071	37,520	9,281	1,934	388	92	155	170,757	1,074	171,831	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	11	167	736	1,090	590	164	24	7	3	5	2,797	3,581	6,378	2.44	1.14
(%)	(0.4)	(6.0)	(26.3)	(39.0)	(21.1)	(5.9)	(0.9)	(0.3)	(0.1)	(0.2)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	10.60	10.53	10.62	10.71	10.79	10.84	10.92	10.76	10.68	10.79	10.71	10.52	10.71		
標準偏差	1.52	1.40	1.26	1.22	1.23	1.27	1.35	1.48	1.38	1.26	1.25	1.54	1.25		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査