

表 2629 体重減少量(kg) 総コレステロール濃度(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

総コレステロール濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
100未満	37	296	1,193	2,079	1,047	266	59	10	2	7	4,996	37	5,033	2.44	1.17
(%)	(0.7)	(5.9)	(23.9)	(41.6)	(21.0)	(5.3)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
100～	71	684	3,177	5,995	3,322	848	161	34	9	9	14,310	120	14,430	2.52	1.04
(%)	(0.5)	(4.8)	(22.2)	(41.9)	(23.2)	(5.9)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
120～	105	1,163	5,782	11,241	6,166	1,498	306	64	15	21	26,361	195	26,556	2.53	1.04
(%)	(0.4)	(4.4)	(21.9)	(42.6)	(23.4)	(5.7)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
140～	90	1,322	6,888	12,676	6,893	1,692	375	74	16	29	30,055	198	30,253	2.53	1.04
(%)	(0.3)	(4.4)	(22.9)	(42.2)	(22.9)	(5.6)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
160～	101	1,063	5,939	10,428	5,432	1,331	266	53	9	32	24,654	127	24,781	2.50	1.05
(%)	(0.4)	(4.3)	(24.1)	(42.3)	(22.0)	(5.4)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
180～	75	755	3,677	6,767	3,191	751	156	29	8	9	15,418	95	15,513	2.45	1.01
(%)	(0.5)	(4.9)	(23.8)	(43.9)	(20.7)	(4.9)	(1.0)	(0.2)	(0.1)	(0.1)	(100.0)				
200～	34	471	2,032	3,421	1,517	400	70	23	1	4	7,973	48	8,021	2.40	1.00
(%)	(0.4)	(5.9)	(25.5)	(42.9)	(19.0)	(5.0)	(0.9)	(0.3)	(0.0)	(0.1)	(100.0)				
220～	20	204	934	1,491	634	135	31	2	1	1	3,453	22	3,475	2.34	0.99
(%)	(0.6)	(5.9)	(27.0)	(43.2)	(18.4)	(3.9)	(0.9)	(0.1)	(0.0)	(0.0)	(100.0)				
240～	9	100	355	563	216	55	7	1			1,306	8	1,314	2.26	0.96
(%)	(0.7)	(7.7)	(27.2)	(43.1)	(16.5)	(4.2)	(0.5)	(0.1)			(100.0)				
260～	14	77	220	311	141	32	8	2		1	806	6	812	2.22	1.12
(%)	(1.7)	(9.6)	(27.3)	(38.6)	(17.5)	(4.0)	(1.0)	(0.2)		(0.1)	(100.0)				
合計	556	6,135	30,197	54,972	28,559	7,008	1,439	292	61	113	129,332	856	130,188	2.49	1.04
(%)	(0.4)	(4.7)	(23.3)	(42.5)	(22.1)	(5.4)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし	199	2,315	10,828	18,189	9,551	2,437	519	103	34	47	44,222	3,799	48,021	2.47	1.08
(%)	(0.5)	(5.2)	(24.5)	(41.1)	(21.6)	(5.5)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	157.35	157.02	156.42	155.06	152.92	152.54	152.18	152.14	146.87	151.42	154.83	151.03	154.80		
標準偏差	44.17	39.27	36.03	34.95	33.68	33.40	33.55	34.90	30.26	31.99	35.12	35.67	35.12		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査