

表 2630 体重減少量(kg) HDL-C濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

HDL-C濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
20未満	7	95	148	139	85	21	7	1			503	11	514	2.04	1.18
(%)	(1.4)	(18.9)	(29.4)	(27.6)	(16.9)	(4.2)	(1.4)	(0.2)			(100.0)				
20～	46	520	2,164	3,869	2,254	717	215	43	14	14	9,856	77	9,933	2.59	1.18
(%)	(0.5)	(5.3)	(22.0)	(39.3)	(22.9)	(7.3)	(2.2)	(0.4)	(0.1)	(0.1)	(100.0)				
30～	112	1,288	6,302	11,586	6,567	1,879	408	90	19	21	28,272	206	28,478	2.56	1.07
(%)	(0.4)	(4.6)	(22.3)	(41.0)	(23.2)	(6.6)	(1.4)	(0.3)	(0.1)	(0.1)	(100.0)				
40～	103	1,347	7,337	13,613	7,053	1,694	315	61	15	39	31,577	201	31,778	2.51	1.05
(%)	(0.3)	(4.3)	(23.2)	(43.1)	(22.3)	(5.4)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
50～	100	997	5,473	9,858	5,007	1,091	188	46	4	14	22,778	127	22,905	2.47	0.99
(%)	(0.4)	(4.4)	(24.0)	(43.3)	(22.0)	(4.8)	(0.8)	(0.2)	(0.0)	(0.1)	(100.0)				
60～	61	550	3,061	5,856	2,921	617	105	20	6	6	13,203	75	13,278	2.47	0.98
(%)	(0.5)	(4.2)	(23.2)	(44.4)	(22.1)	(4.7)	(0.8)	(0.2)	(0.0)	(0.0)	(100.0)				
70～	30	317	1,524	2,873	1,418	290	66	6	3	8	6,535	40	6,575	2.46	1.03
(%)	(0.5)	(4.9)	(23.3)	(44.0)	(21.7)	(4.4)	(1.0)	(0.1)	(0.0)	(0.1)	(100.0)				
80～	13	142	753	1,376	644	144	33	7			3,112	13	3,125	2.45	0.95
(%)	(0.4)	(4.6)	(24.2)	(44.2)	(20.7)	(4.6)	(1.1)	(0.2)			(100.0)				
90～	9	81	343	583	297	65	15	3		1	1,397	11	1,408	2.45	1.06
(%)	(0.6)	(5.8)	(24.6)	(41.7)	(21.3)	(4.7)	(1.1)	(0.2)		(0.1)	(100.0)				
100～	9	61	264	503	248	57	7	2		1	1,152	9	1,161	2.43	1.00
(%)	(0.8)	(5.3)	(22.9)	(43.7)	(21.5)	(4.9)	(0.6)	(0.2)		(0.1)	(100.0)				
合計	490	5,398	27,369	50,256	26,494	6,575	1,359	279	61	104	118,385	770	119,155	2.51	1.04
(%)	(0.4)	(4.6)	(23.1)	(42.5)	(22.4)	(5.6)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	265	3,052	13,656	22,905	11,616	2,870	599	116	34	56	55,169	3,885	59,054	2.44	1.06
(%)	(0.5)	(5.5)	(24.8)	(41.5)	(21.1)	(5.2)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	49.57	47.92	48.78	48.98	48.11	46.22	44.44	43.83	41.07	45.38	48.47	46.90	48.46		
標準偏差	19.34	17.67	16.58	16.59	16.39	16.25	16.39	16.43	13.99	16.06	16.61	18.01	16.62		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査