

表 2631 体重減少量(kg) Non-HDL-C濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

Non-HDL-C濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
50未満	20	128	478	894	477	114	24	4	2	1	2,142	15	2,157	2.46	1.05
(%)	(0.9)	(6.0)	(22.3)	(41.7)	(22.3)	(5.3)	(1.1)	(0.2)	(0.1)	(0.0)	(100.0)				
50～	50	408	2,117	4,248	2,431	628	132	31	5	7	10,057	77	10,134	2.58	1.10
(%)	(0.5)	(4.1)	(21.1)	(42.2)	(24.2)	(6.2)	(1.3)	(0.3)	(0.0)	(0.1)	(100.0)				
70～	86	921	4,854	9,443	5,248	1,228	228	46	8	17	22,079	161	22,240	2.53	1.01
(%)	(0.4)	(4.2)	(22.0)	(42.8)	(23.8)	(5.6)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
90～	83	1,113	6,108	11,211	5,768	1,335	275	54	14	25	25,986	191	26,177	2.50	1.02
(%)	(0.3)	(4.3)	(23.5)	(43.1)	(22.2)	(5.1)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
110～	76	1,010	4,985	8,909	4,397	1,135	241	43	15	21	20,832	151	20,983	2.48	1.05
(%)	(0.4)	(4.8)	(23.9)	(42.8)	(21.1)	(5.4)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
130～	61	592	3,012	5,327	2,713	676	139	37	4	11	12,572	62	12,634	2.48	1.07
(%)	(0.5)	(4.7)	(24.0)	(42.4)	(21.6)	(5.4)	(1.1)	(0.3)	(0.0)	(0.1)	(100.0)				
150～	25	359	1,529	2,604	1,299	339	92	14	4	5	6,270	39	6,309	2.46	1.04
(%)	(0.4)	(5.7)	(24.4)	(41.5)	(20.7)	(5.4)	(1.5)	(0.2)	(0.1)	(0.1)	(100.0)				
170～	12	154	676	1,120	551	169	32	11	2	1	2,728	27	2,755	2.45	1.05
(%)	(0.4)	(5.6)	(24.8)	(41.1)	(20.2)	(6.2)	(1.2)	(0.4)	(0.1)	(0.0)	(100.0)				
190～	11	63	252	423	203	53	11	2			1,018	6	1,024	2.37	1.02
(%)	(1.1)	(6.2)	(24.8)	(41.6)	(19.9)	(5.2)	(1.1)	(0.2)			(100.0)				
210～	4	52	141	228	111	35	10	2		1	584	5	589	2.40	1.16
(%)	(0.7)	(8.9)	(24.1)	(39.0)	(19.0)	(6.0)	(1.7)	(0.3)		(0.2)	(100.0)				
合計	428	4,800	24,152	44,407	23,198	5,712	1,184	244	54	89	104,268	734	105,002	2.50	1.04
(%)	(0.4)	(4.6)	(23.2)	(42.6)	(22.2)	(5.5)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
記載なし	327	3,650	16,873	28,754	14,912	3,733	774	151	41	71	69,286	3,921	73,207	2.46	1.06
(%)	(0.5)	(5.3)	(24.4)	(41.5)	(21.5)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	107.07	109.54	107.69	106.09	104.76	105.94	107.73	108.28	107.24	106.55	106.35	104.99	106.34		
標準偏差	39.40	35.81	33.21	32.74	32.80	33.62	35.50	36.08	32.07	31.26	33.15	33.02	33.15		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査