

表 2632 体重減少量(kg) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前収縮期血圧	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	1	3	7	10	1	3					25	1	26	2.04	1.23
(%)	(4.0)	(12.0)	(28.0)	(40.0)	(4.0)	(12.0)					(100.0)				
60～	10	41	68	80	28	1	2				230	3	233	1.84	1.08
(%)	(4.3)	(17.8)	(29.6)	(34.8)	(12.2)	(0.4)	(0.9)				(100.0)				
80～	40	215	652	711	228	53	12	1			1,912	11	1,923	2.00	1.00
(%)	(2.1)	(11.2)	(34.1)	(37.2)	(11.9)	(2.8)	(0.6)	(0.1)			(100.0)				
100～	102	834	3,135	4,034	1,498	307	64	9	2	10	9,995	48	10,043	2.19	1.05
(%)	(1.0)	(8.3)	(31.4)	(40.4)	(15.0)	(3.1)	(0.6)	(0.1)	(0.0)	(0.1)	(100.0)				
120～	166	1,854	8,498	13,509	6,008	1,265	232	47	12	30	31,621	129	31,750	2.36	1.01
(%)	(0.5)	(5.9)	(26.9)	(42.7)	(19.0)	(4.0)	(0.7)	(0.1)	(0.0)	(0.1)	(100.0)				
140～	194	2,223	11,404	21,736	11,161	2,418	468	71	17	47	49,739	159	49,898	2.49	1.01
(%)	(0.4)	(4.5)	(22.9)	(43.7)	(22.4)	(4.9)	(0.9)	(0.1)	(0.0)	(0.1)	(100.0)				
160～	99	1,396	7,797	15,479	9,063	2,348	477	111	19	28	36,817	125	36,942	2.59	1.04
(%)	(0.3)	(3.8)	(21.2)	(42.0)	(24.6)	(6.4)	(1.3)	(0.3)	(0.1)	(0.1)	(100.0)				
180～	38	485	2,705	5,693	3,687	1,205	263	62	14	15	14,167	51	14,218	2.70	1.11
(%)	(0.3)	(3.4)	(19.1)	(40.2)	(26.0)	(8.5)	(1.9)	(0.4)	(0.1)	(0.1)	(100.0)				
200～	8	108	595	1,347	1,040	361	99	25	3	2	3,588	15	3,603	2.84	1.14
(%)	(0.2)	(3.0)	(16.6)	(37.5)	(29.0)	(10.1)	(2.8)	(0.7)	(0.1)	(0.1)	(100.0)				
220～	1	13	135	262	193	82	24	8		2	720	3	723	2.88	1.21
(%)	(0.1)	(1.8)	(18.8)	(36.4)	(26.8)	(11.4)	(3.3)	(1.1)		(0.3)	(100.0)				
合計	659	7,172	34,996	62,861	32,907	8,043	1,641	334	67	134	148,814	545	149,359	2.49	1.05
(%)	(0.4)	(4.8)	(23.5)	(42.2)	(22.1)	(5.4)	(1.1)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし	96	1,278	6,029	10,300	5,203	1,402	317	61	28	26	24,740	4,110	28,850	2.48	1.08
(%)	(0.4)	(5.2)	(24.4)	(41.6)	(21.0)	(5.7)	(1.3)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	755	8,450	41,025	73,161	38,110	9,445	1,958	395	95	160	173,554	4,655	178,209	2.49	1.05
(%)	(0.4)	(4.9)	(23.6)	(42.2)	(22.0)	(5.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	139.69	144.76	148.31	151.68	155.53	159.30	161.32	165.70	161.49	151.84	151.91	149.28	151.90		
標準偏差	27.56	25.30	24.31	23.40	23.37	24.53	25.72	25.47	24.71	24.29	24.10	26.48	24.11		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査