

表 2638 体重減少量(kg) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	42	488	3,136	6,181	3,116	698	119	19	4	7	13,810	316	14,126	2.51	0.96
(%)	(0.3)	(3.5)	(22.7)	(44.8)	(22.6)	(5.1)	(0.9)	(0.1)	(0.0)	(0.1)	(100.0)				
慢性腎盂腎炎	2	20	104	162	74	23	4				389	9	398	2.42	0.99
(%)	(0.5)	(5.1)	(26.7)	(41.6)	(19.0)	(5.9)	(1.0)				(100.0)				
急速進行性糸球体腎炎	1	10	42	122	48	20	2				245	5	250	2.53	0.93
(%)	(0.4)	(4.1)	(17.1)	(49.8)	(19.6)	(8.2)	(0.8)				(100.0)				
妊娠腎/妊娠中毒症	2	10	109	179	41	1					342	6	348	2.19	0.72
(%)	(0.6)	(2.9)	(31.9)	(52.3)	(12.0)	(0.3)					(100.0)				
その他の分類不能の腎炎	2	3	43	69	42	19	3	1			182	4	186	2.67	1.05
(%)	(1.1)	(1.6)	(23.6)	(37.9)	(23.1)	(10.4)	(1.6)	(0.5)			(100.0)				
多発性嚢胞腎	2	52	318	579	283	54	5	1		5	1,299	31	1,330	2.48	1.04
(%)	(0.2)	(4.0)	(24.5)	(44.6)	(21.8)	(4.2)	(0.4)	(0.1)		(0.4)	(100.0)				
腎硬化症	5	107	564	916	467	139	19	5			2,222	30	2,252	2.47	0.98
(%)	(0.2)	(4.8)	(25.4)	(41.2)	(21.0)	(6.3)	(0.9)	(0.2)			(100.0)				
悪性高血圧	1	8	35	99	85	32	5				265	5	270	2.87	1.00
(%)	(0.4)	(3.0)	(13.2)	(37.4)	(32.1)	(12.1)	(1.9)				(100.0)				
糖尿病性腎症	23	395	1,888	4,291	2,825	930	186	23	10	10	10,581	264	10,845	2.71	1.10
(%)	(0.2)	(3.7)	(17.8)	(40.6)	(26.7)	(8.8)	(1.8)	(0.2)	(0.1)	(0.1)	(100.0)				
SLE腎炎	1	25	92	149	48	13	3			1	332	5	337	2.37	1.42
(%)	(0.3)	(7.5)	(27.7)	(44.9)	(14.5)	(3.9)	(0.9)			(0.3)	(100.0)				
アミロイド腎		3	10	19	10	1					43		43	2.36	0.84
(%)		(7.0)	(23.3)	(44.2)	(23.3)	(2.3)					(100.0)				
痛風腎		5	27	64	34	6					136	6	142	2.50	0.90
(%)		(3.7)	(19.9)	(47.1)	(25.0)	(4.4)					(100.0)				
先天性代謝異常による腎不全		1	8	17	7	6					39	2	41	2.69	1.05
(%)		(2.6)	(20.5)	(43.6)	(17.9)	(15.4)					(100.0)				
腎・尿路結核		2	8	8	2						20		20	2.03	0.73
(%)		(10.0)	(40.0)	(40.0)	(10.0)						(100.0)				
腎・尿路結石		2	17	25	16	3					63	1	64	2.43	0.94
(%)		(3.2)	(27.0)	(39.7)	(25.4)	(4.8)					(100.0)				
腎・尿路腫瘍		4	9	32	14	2	1				62		62	2.49	0.97
(%)		(6.5)	(14.5)	(51.6)	(22.6)	(3.2)	(1.6)				(100.0)				
閉塞性尿路障害		3	17	23	18	4		1			66	3	69	2.58	1.05
(%)		(4.5)	(25.8)	(34.8)	(27.3)	(6.1)		(1.5)			(100.0)				
骨髄腫		1	4	3	3						11		11	2.15	0.98
(%)		(9.1)	(36.4)	(27.3)	(27.3)						(100.0)				
腎形成不全		3	22	44	20	5					94	1	95	2.48	0.87
(%)		(3.2)	(23.4)	(46.8)	(21.3)	(5.3)					(100.0)				
不明	12	141	698	1,206	598	166	35	5		5	2,866	88	2,954	2.49	1.07
(%)	(0.4)	(4.9)	(24.4)	(42.1)	(20.9)	(5.8)	(1.2)	(0.2)		(0.2)	(100.0)				
再導入	1	3	74	172	94	28	4		1		377	8	385	2.71	0.93
(%)	(0.3)	(0.8)	(19.6)	(45.6)	(24.9)	(7.4)	(1.1)		(0.3)		(100.0)				
その他	3	34	175	294	169	41	14	4	1	2	737	6	743	2.56	1.18
(%)	(0.4)	(4.6)	(23.7)	(39.9)	(22.9)	(5.6)	(1.9)	(0.5)	(0.1)	(0.3)	(100.0)				
合計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし															
(%)															
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査