

表 2639 体重減少量(kg) 医療機関別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

医療機関	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	1	5	11	13	6	1					37		37	2.06	1.06
(%)	(2.7)	(13.5)	(29.7)	(35.1)	(16.2)	(2.7)					(100.0)				
私立大学	2	4	48	58	28	6	2	1			149	2	151	2.40	0.98
(%)	(1.3)	(2.7)	(32.2)	(38.9)	(18.8)	(4.0)	(1.3)	(0.7)			(100.0)				
国立		3	13	7	7	2					32		32	2.23	1.12
(%)		(9.4)	(40.6)	(21.9)	(21.9)	(6.3)					(100.0)				
県市町村立	4	58	331	681	341	78	9	4			1,506	33	1,539	2.51	0.93
(%)	(0.3)	(3.9)	(22.0)	(45.2)	(22.6)	(5.2)	(0.6)	(0.3)			(100.0)				
社会保険		16	56	141	53	15	1				282	9	291	2.45	0.91
(%)		(5.7)	(19.9)	(50.0)	(18.8)	(5.3)	(0.4)				(100.0)				
厚生連	1	21	171	371	226	59	13	1	1	1	865	40	905	2.67	1.01
(%)	(0.1)	(2.4)	(19.8)	(42.9)	(26.1)	(6.8)	(1.5)	(0.1)	(0.1)	(0.1)	(100.0)				
その他公的	2	30	133	238	141	37	4	4		2	591	27	618	2.57	1.28
(%)	(0.3)	(5.1)	(22.5)	(40.3)	(23.9)	(6.3)	(0.7)	(0.7)		(0.3)	(100.0)				
私立総合	4	29	199	441	205	48	8	1	1	1	937	18	955	2.53	0.94
(%)	(0.4)	(3.1)	(21.2)	(47.1)	(21.9)	(5.1)	(0.9)	(0.1)	(0.1)	(0.1)	(100.0)				
私立	23	343	1,942	3,575	1,951	557	88	9	5	5	8,498	328	8,826	2.55	1.05
(%)	(0.3)	(4.0)	(22.9)	(42.1)	(23.0)	(6.6)	(1.0)	(0.1)	(0.1)	(0.1)	(100.0)				
私立診療所	60	811	4,496	9,129	5,056	1,388	275	39	9	21	21,284	333	21,617	2.58	1.03
(%)	(0.3)	(3.8)	(21.1)	(42.9)	(23.8)	(6.5)	(1.3)	(0.2)	(0.0)	(0.1)	(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査