

表 2646 体重減少量(kg) Body Mass Index別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

Body Mass Index	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
14未満	4	25	57	36	3					2	127	11	138	1.75	2.14
(%)	(3.1)	(19.7)	(44.9)	(28.3)	(2.4)					(1.6)	(100.0)				
14～	7	92	422	358	70	11	1			2	963	1	964	1.92	0.92
(%)	(0.7)	(9.6)	(43.8)	(37.2)	(7.3)	(1.1)	(0.1)			(0.2)	(100.0)				
16～	10	214	1,173	1,737	462	53	2	1		5	3,657	2	3,659	2.18	0.93
(%)	(0.3)	(5.9)	(32.1)	(47.5)	(12.6)	(1.4)	(0.1)	(0.0)		(0.1)	(100.0)				
18～	17	286	1,818	3,240	1,255	201	16	5	2	8	6,848	1	6,849	2.36	0.91
(%)	(0.2)	(4.2)	(26.5)	(47.3)	(18.3)	(2.9)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
20～	28	271	1,603	3,388	1,752	335	39	6	3	7	7,432		7,432	2.52	0.99
(%)	(0.4)	(3.6)	(21.6)	(45.6)	(23.6)	(4.5)	(0.5)	(0.1)	(0.0)	(0.1)	(100.0)				
22～	10	182	1,032	2,480	1,595	404	48	7	3		5,761	1	5,762	2.66	0.94
(%)	(0.2)	(3.2)	(17.9)	(43.0)	(27.7)	(7.0)	(0.8)	(0.1)	(0.1)		(100.0)				
24～	13	88	492	1,458	1,098	352	57	4			3,562		3,562	2.81	0.98
(%)	(0.4)	(2.5)	(13.8)	(40.9)	(30.8)	(9.9)	(1.6)	(0.1)			(100.0)				
26～	3	35	207	643	632	267	61	6		1	1,855		1,855	3.04	1.04
(%)	(0.2)	(1.9)	(11.2)	(34.7)	(34.1)	(14.4)	(3.3)	(0.3)		(0.1)	(100.0)				
28～	2	15	95	306	349	181	47	3			998		998	3.19	1.04
(%)	(0.2)	(1.5)	(9.5)	(30.7)	(35.0)	(18.1)	(4.7)	(0.3)			(100.0)				
30～		17	60	186	289	203	62	13	3	1	834	2	836	3.48	1.19
(%)		(2.0)	(7.2)	(22.3)	(34.7)	(24.3)	(7.4)	(1.6)	(0.4)	(0.1)	(100.0)				
合計	94	1,225	6,959	13,832	7,505	2,007	333	45	11	26	32,037	18	32,055	2.56	1.02
(%)	(0.3)	(3.8)	(21.7)	(43.2)	(23.4)	(6.3)	(1.0)	(0.1)	(0.0)	(0.1)	(100.0)				
記載なし	3	95	441	822	509	184	67	14	5	4	2,144	772	2,916	2.71	1.18
(%)	(0.1)	(4.4)	(20.6)	(38.3)	(23.7)	(8.6)	(3.1)	(0.7)	(0.2)	(0.2)	(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	20.55	20.21	20.33	21.23	22.65	24.40	26.19	25.98	24.20	19.27	21.58	13.29	21.58		
標準偏差	3.48	3.45	3.22	3.27	3.57	3.91	4.11	4.96	5.49	3.73	3.61	8.07	3.62		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査