

表 2647 体重減少量(kg) Body Mass Index別（血液透析濾過患者 男性のみ 透析回数週3回 透析歴2年以上）

Body Mass Index	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
14未満	1	8	12	11	1					2	35	5	40	2.34	3.85
(%)	(2.9)	(22.9)	(34.3)	(31.4)	(2.9)					(5.7)	(100.0)				
14～	1	30	111	135	40	6					323	1	324	2.05	0.85
(%)	(0.3)	(9.3)	(34.4)	(41.8)	(12.4)	(1.9)					(100.0)				
16～	4	76	430	795	307	41	2	1		3	1,659	1	1,660	2.36	1.01
(%)	(0.2)	(4.6)	(25.9)	(47.9)	(18.5)	(2.5)	(0.1)	(0.1)		(0.2)	(100.0)				
18～	8	136	796	1,802	950	189	16	5	1	5	3,908		3,908	2.54	0.95
(%)	(0.2)	(3.5)	(20.4)	(46.1)	(24.3)	(4.8)	(0.4)	(0.1)	(0.0)	(0.1)	(100.0)				
20～	20	149	878	2,109	1,398	308	36	6	2	6	4,912		4,912	2.66	1.05
(%)	(0.4)	(3.0)	(17.9)	(42.9)	(28.5)	(6.3)	(0.7)	(0.1)	(0.0)	(0.1)	(100.0)				
22～	8	119	574	1,613	1,318	373	46	4	3		4,058	1	4,059	2.80	0.96
(%)	(0.2)	(2.9)	(14.1)	(39.7)	(32.5)	(9.2)	(1.1)	(0.1)	(0.1)		(100.0)				
24～	11	55	268	917	879	325	54	3			2,512		2,512	2.97	1.00
(%)	(0.4)	(2.2)	(10.7)	(36.5)	(35.0)	(12.9)	(2.1)	(0.1)			(100.0)				
26～	3	22	107	379	490	246	61	6		1	1,315		1,315	3.24	1.06
(%)	(0.2)	(1.7)	(8.1)	(28.8)	(37.3)	(18.7)	(4.6)	(0.5)		(0.1)	(100.0)				
28～	2	9	45	162	261	160	42	3			684		684	3.41	1.04
(%)	(0.3)	(1.3)	(6.6)	(23.7)	(38.2)	(23.4)	(6.1)	(0.4)			(100.0)				
30～		10	20	85	195	170	56	11	2	1	550	2	552	3.76	1.15
(%)		(1.8)	(3.6)	(15.5)	(35.5)	(30.9)	(10.2)	(2.0)	(0.4)	(0.2)	(100.0)				
合計	58	614	3,241	8,008	5,839	1,818	313	39	8	18	19,956	10	19,966	2.76	1.06
(%)	(0.3)	(3.1)	(16.2)	(40.1)	(29.3)	(9.1)	(1.6)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし		47	200	442	384	160	62	13	5	4	1,317	493	1,810	2.99	1.25
(%)		(3.6)	(15.2)	(33.6)	(29.2)	(12.1)	(4.7)	(1.0)	(0.4)	(0.3)	(100.0)				
総計	58	661	3,441	8,450	6,223	1,978	375	52	13	22	21,273	503	21,776	2.78	1.08
(%)	(0.3)	(3.1)	(16.2)	(39.7)	(29.3)	(9.3)	(1.8)	(0.2)	(0.1)	(0.1)	(100.0)				
平均	21.82	20.88	20.80	21.46	22.68	24.36	26.16	25.98	24.47	19.92	22.04	15.71	22.04		
標準偏差	3.08	3.36	3.00	3.08	3.42	3.82	4.06	5.08	5.27	4.13	3.47	9.34	3.48		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査