

表 2648 体重減少量(kg) Body Mass Index別 (血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

Body Mass Index	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
14未満	3	17	45	25	2						92	6	98	1.53	0.77
(%)	(3.3)	(18.5)	(48.9)	(27.2)	(2.2)						(100.0)				
14～	6	62	311	223	30	5	1			2	640		640	1.85	0.95
(%)	(0.9)	(9.7)	(48.6)	(34.8)	(4.7)	(0.8)	(0.2)			(0.3)	(100.0)				
16～	6	138	743	942	155	12				2	1,998	1	1,999	2.03	0.82
(%)	(0.3)	(6.9)	(37.2)	(47.1)	(7.8)	(0.6)				(0.1)	(100.0)				
18～	9	150	1,022	1,438	305	12			1	3	2,940	1	2,941	2.13	0.79
(%)	(0.3)	(5.1)	(34.8)	(48.9)	(10.4)	(0.4)			(0.0)	(0.1)	(100.0)				
20～	8	122	725	1,279	354	27	3			1	2,520		2,520	2.23	0.78
(%)	(0.3)	(4.8)	(28.8)	(50.8)	(14.0)	(1.1)	(0.1)		(0.0)	(0.0)	(100.0)				
22～	2	63	458	867	277	31	2	3			1,703		1,703	2.31	0.78
(%)	(0.1)	(3.7)	(26.9)	(50.9)	(16.3)	(1.8)	(0.1)	(0.2)			(100.0)				
24～	2	33	224	541	219	27	3	1			1,050		1,050	2.44	0.81
(%)	(0.2)	(3.1)	(21.3)	(51.5)	(20.9)	(2.6)	(0.3)	(0.1)			(100.0)				
26～		13	100	264	142	21					540		540	2.55	0.81
(%)		(2.4)	(18.5)	(48.9)	(26.3)	(3.9)					(100.0)				
28～		6	50	144	88	21	5				314		314	2.71	0.86
(%)		(1.9)	(15.9)	(45.9)	(28.0)	(6.7)	(1.6)				(100.0)				
30～		7	40	101	94	33	6	2	1		284		284	2.94	1.06
(%)		(2.5)	(14.1)	(35.6)	(33.1)	(11.6)	(2.1)	(0.7)	(0.4)		(100.0)				
合計	36	611	3,718	5,824	1,666	189	20	6	3	8	12,081	8	12,089	2.22	0.84
(%)	(0.3)	(5.1)	(30.8)	(48.2)	(13.8)	(1.6)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
記載なし	3	48	241	380	125	24	5	1			827	279	1,106	2.27	0.88
(%)	(0.4)	(5.8)	(29.1)	(45.9)	(15.1)	(2.9)	(0.6)	(0.1)			(100.0)				
総計	39	659	3,959	6,204	1,791	213	25	7	3	8	12,908	287	13,195	2.22	0.84
(%)	(0.3)	(5.1)	(30.7)	(48.1)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
平均	18.51	19.54	19.93	20.91	22.54	24.83	26.77	25.99	23.46	17.80	20.83	10.25	20.82		
標準偏差	3.12	3.41	3.34	3.50	4.05	4.69	4.97	4.55	7.24	2.18	3.70	5.21	3.71		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査