

表 2649 体重減少量(kg) 透析前BUN(mg/dL)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前BUN	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	14	11	7	3						37	2	39	1.28	1.05
(%)	(5.4)	(37.8)	(29.7)	(18.9)	(8.1)						(100.0)				
20～	7	68	119	78	23	4	2				301	8	309	1.68	1.01
(%)	(2.3)	(22.6)	(39.5)	(25.9)	(7.6)	(1.3)	(0.7)				(100.0)				
30～	19	168	571	617	204	51	8	3	1		1,642	11	1,653	2.08	0.99
(%)	(1.2)	(10.2)	(34.8)	(37.6)	(12.4)	(3.1)	(0.5)	(0.2)	(0.1)		(100.0)				
40～	19	288	1,362	2,023	855	169	27	3		7	4,753	32	4,785	2.32	1.00
(%)	(0.4)	(6.1)	(28.7)	(42.6)	(18.0)	(3.6)	(0.6)	(0.1)		(0.1)	(100.0)				
50～	20	317	1,935	3,650	1,816	438	75	8	3	10	8,272	46	8,318	2.50	1.02
(%)	(0.2)	(3.8)	(23.4)	(44.1)	(22.0)	(5.3)	(0.9)	(0.1)	(0.0)	(0.1)	(100.0)				
60～	18	231	1,752	3,915	2,151	560	100	21	6	8	8,762	42	8,804	2.63	0.98
(%)	(0.2)	(2.6)	(20.0)	(44.7)	(24.5)	(6.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
70～	9	140	1,001	2,638	1,682	494	73	13	3	2	6,055	31	6,086	2.74	1.01
(%)	(0.1)	(2.3)	(16.5)	(43.6)	(27.8)	(8.2)	(1.2)	(0.2)	(0.0)	(0.0)	(100.0)				
80～	3	52	417	1,145	851	286	59	2	1	1	2,817	14	2,831	2.85	1.00
(%)	(0.1)	(1.8)	(14.8)	(40.6)	(30.2)	(10.2)	(2.1)	(0.1)	(0.0)	(0.0)	(100.0)				
90～		19	131	401	319	127	41	6	1	1	1,046	5	1,051	2.99	1.08
(%)		(1.8)	(12.5)	(38.3)	(30.5)	(12.1)	(3.9)	(0.6)	(0.1)	(0.1)	(100.0)				
100～		3	40	106	68	46	9	3	1		276	3	279	3.04	1.15
(%)		(1.1)	(14.5)	(38.4)	(24.6)	(16.7)	(3.3)	(1.1)	(0.4)		(100.0)				
110～		1	6	25	23	14	5				74		74	3.19	1.12
(%)		(1.4)	(8.1)	(33.8)	(31.1)	(18.9)	(6.8)				(100.0)				
120～			3	11	4	1	1				20		20	2.65	0.98
(%)			(15.0)	(55.0)	(20.0)	(5.0)	(5.0)				(100.0)				
合計	97	1,301	7,348	14,616	7,999	2,190	400	59	16	29	34,055	194	34,249	2.57	1.03
(%)	(0.3)	(3.8)	(21.6)	(42.9)	(23.5)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし		19	52	38	15	1				1	126	596	722	1.94	1.05
(%)		(15.1)	(41.3)	(30.2)	(11.9)	(0.8)				(0.8)	(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	50.47	53.57	58.41	62.37	65.10	68.05	70.01	68.61	67.63	59.28	62.25	59.10	62.23		
標準偏差	15.39	16.10	14.88	14.41	14.52	15.26	16.48	16.52	16.46	11.88	15.07	17.26	15.08		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査