

表 2653 体重減少量(kg) single pool Kt/V(Kt/Vsp)別(血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

single pool Kt/V (Kt/Vsp)	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.4未満		4	4	1	1						10		10	1.37	0.94
(%)		(40.0)	(40.0)	(10.0)	(10.0)						(100.0)				
0.4～			4	1	1						6		6	1.88	0.67
(%)			(66.7)	(16.7)	(16.7)						(100.0)				
0.6～		5	8	6							19		19	1.39	0.70
(%)		(26.3)	(42.1)	(31.6)							(100.0)				
0.8～	1	15	36	22	11						85		85	1.79	0.90
(%)	(1.2)	(17.6)	(42.4)	(25.9)	(12.9)						(100.0)				
1.0～	6	49	118	160	43	15	4				395		395	2.08	1.04
(%)	(1.5)	(12.4)	(29.9)	(40.5)	(10.9)	(3.8)	(1.0)				(100.0)				
1.2～	7	92	431	584	208	33	3	1			1,359		1,359	2.20	0.89
(%)	(0.5)	(6.8)	(31.7)	(43.0)	(15.3)	(2.4)	(0.2)	(0.1)			(100.0)				
1.4～	11	149	831	1,298	426	54	7	2			2,778		2,778	2.24	0.83
(%)	(0.4)	(5.4)	(29.9)	(46.7)	(15.3)	(1.9)	(0.3)	(0.1)			(100.0)				
1.6～	9	150	1,007	1,733	469	48	5	2			3,423		3,423	2.23	0.76
(%)	(0.3)	(4.4)	(29.4)	(50.6)	(13.7)	(1.4)	(0.1)	(0.1)			(100.0)				
1.8～	3	83	737	1,217	320	26	1	1			2,388		2,388	2.23	0.72
(%)	(0.1)	(3.5)	(30.9)	(51.0)	(13.4)	(1.1)	(0.0)	(0.0)			(100.0)				
2.0～	1	67	587	957	247	32	4		2	7	1,904		1,904	2.29	1.00
(%)	(0.1)	(3.5)	(30.8)	(50.3)	(13.0)	(1.7)	(0.2)		(0.1)	(0.4)	(100.0)				
合計	38	614	3,763	5,979	1,726	208	24	6	2	7	12,367		12,367	2.23	0.84
(%)	(0.3)	(5.0)	(30.4)	(48.3)	(14.0)	(1.7)	(0.2)	(0.0)	(0.0)	(0.1)	(100.0)				
記載なし	1	45	196	225	65	5	1	1	1	1	541	287	828	2.10	0.96
(%)	(0.2)	(8.3)	(36.2)	(41.6)	(12.0)	(0.9)	(0.2)	(0.2)	(0.2)	(0.2)	(100.0)				
総計	39	659	3,959	6,204	1,791	213	25	7	3	8	12,908	287	13,195	2.22	0.84
(%)	(0.3)	(5.1)	(30.7)	(48.1)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
平均	1.46	1.58	1.69	1.72	1.69	1.64	1.64	1.63	2.58	2.52	1.70		1.70		
標準偏差	0.29	0.35	0.32	0.31	0.31	0.33	0.44	0.21	0.37	0.20	0.32		0.32		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査