

表 2656 体重減少量(kg) equilibrated Kt/V(eKt/V)別(血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

equilibrated Kt/V (eKt/V)	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
0.4未満		4	5	1	1						11		11	1.35	0.90
(%)		(36.4)	(45.5)	(9.1)	(9.1)						(100.0)				
0.4～		2	5	4	1						12		12	1.68	0.79
(%)		(16.7)	(41.7)	(33.3)	(8.3)						(100.0)				
0.6～	1	12	28	17	5						63		63	1.63	0.81
(%)	(1.6)	(19.0)	(44.4)	(27.0)	(7.9)						(100.0)				
0.8～	6	45	109	120	35	8					323		323	1.95	0.97
(%)	(1.9)	(13.9)	(33.7)	(37.2)	(10.8)	(2.5)					(100.0)				
1.0～	8	103	452	588	191	37	6		1		1,386		1,386	2.17	0.91
(%)	(0.6)	(7.4)	(32.6)	(42.4)	(13.8)	(2.7)	(0.4)		(0.1)		(100.0)				
1.2～	10	147	869	1,453	487	61	8		2		3,037		3,037	2.27	0.82
(%)	(0.3)	(4.8)	(28.6)	(47.8)	(16.0)	(2.0)	(0.3)		(0.1)		(100.0)				
1.4～	8	156	1,095	1,881	523	49	5		2		3,719		3,719	2.24	0.75
(%)	(0.2)	(4.2)	(29.4)	(50.6)	(14.1)	(1.3)	(0.1)		(0.1)		(100.0)				
1.6～	4	80	710	1,145	295	27	1		1		2,263		2,263	2.22	0.73
(%)	(0.2)	(3.5)	(31.4)	(50.6)	(13.0)	(1.2)	(0.0)		(0.0)		(100.0)				
1.8～		39	297	445	106	19	1			3	910		910	2.23	0.94
(%)		(4.3)	(32.6)	(48.9)	(11.6)	(2.1)	(0.1)			(0.3)	(100.0)				
2.0～	1	26	195	332	82	7	3		1	2	654		655	2.36	1.27
(%)	(0.2)	(4.0)	(29.8)	(50.8)	(12.5)	(1.1)	(0.5)		(0.2)	(0.3)	(100.0)	1	655		
合計	38	614	3,765	5,986	1,726	208	24	7	2	8	12,378	1	12,379	2.23	0.84
(%)	(0.3)	(5.0)	(30.4)	(48.4)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
記載なし	1	45	194	218	65	5	1		1		530	286	816	2.07	0.85
(%)	(0.2)	(8.5)	(36.6)	(41.1)	(12.3)	(0.9)	(0.2)		(0.2)		(100.0)				
総計	39	659	3,959	6,204	1,791	213	25	7	3	8	12,908	287	13,195	2.22	0.84
(%)	(0.3)	(5.1)	(30.7)	(48.1)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
平均	1.28	1.39	1.48	1.50	1.47	1.43	1.44	1.63	2.31	2.16	1.48	2.95	1.48		
標準偏差	0.28	0.34	0.31	0.29	0.28	0.29	0.42	0.55	0.25	0.30	0.30		0.30		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査