

表 2660 体重減少量(kg) %クレアチニン産生速度(%)別 (血液透析濾過患者 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	22	240	589	560	181	37	5				1,634		1,634	1.91	0.98
(%)	(1.3)	(14.7)	(36.0)	(34.3)	(11.1)	(2.3)	(0.3)				(100.0)				
60～	16	133	440	545	208	42	16	1	2	1	1,404		1,404	2.19	1.12
(%)	(1.1)	(9.5)	(31.3)	(38.8)	(14.8)	(3.0)	(1.1)	(0.1)	(0.1)	(0.1)	(100.0)				
70～	15	164	678	1,017	449	117	30	5	1	1	2,477	1	2,478	2.35	1.03
(%)	(0.6)	(6.6)	(27.4)	(41.1)	(18.1)	(4.7)	(1.2)	(0.2)	(0.0)	(0.0)	(100.0)				
80～	14	184	948	1,647	795	234	35	6	3	2	3,868		3,868	2.47	1.00
(%)	(0.4)	(4.8)	(24.5)	(42.6)	(20.6)	(6.0)	(0.9)	(0.2)	(0.1)	(0.1)	(100.0)				
90～	8	163	1,120	2,210	1,185	345	63	5			5,099	1	5,100	2.58	0.95
(%)	(0.2)	(3.2)	(22.0)	(43.3)	(23.2)	(6.8)	(1.2)	(0.1)			(100.0)				
100～	7	130	1,205	2,555	1,449	392	79	11	3	3	5,834		5,834	2.64	0.97
(%)	(0.1)	(2.2)	(20.7)	(43.8)	(24.8)	(6.7)	(1.4)	(0.2)	(0.1)	(0.1)	(100.0)				
110～	5	109	945	2,342	1,522	398	60	11	3	4	5,399		5,399	2.71	0.94
(%)	(0.1)	(2.0)	(17.5)	(43.4)	(28.2)	(7.4)	(1.1)	(0.2)	(0.1)	(0.1)	(100.0)				
120～	3	68	634	1,766	1,025	284	55	6		2	3,843		3,843	2.73	0.96
(%)	(0.1)	(1.8)	(16.5)	(46.0)	(26.7)	(7.4)	(1.4)	(0.2)		(0.1)	(100.0)				
130～	2	31	327	960	564	176	18	9	1	4	2,092		2,092	2.77	0.99
(%)	(0.1)	(1.5)	(15.6)	(45.9)	(27.0)	(8.4)	(0.9)	(0.4)	(0.0)	(0.2)	(100.0)				
140～		13	190	558	391	113	24	2	2	12	1,305	3	1,308	2.94	1.52
(%)		(1.0)	(14.6)	(42.8)	(30.0)	(8.7)	(1.8)	(0.2)	(0.2)	(0.9)	(100.0)				
合計	92	1,235	7,076	14,160	7,769	2,138	385	56	15	29	32,955	5	32,960	2.58	1.03
(%)	(0.3)	(3.7)	(21.5)	(43.0)	(23.6)	(6.5)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし	5	85	324	494	245	53	15	3	1	1	1,226	785	2,011	2.38	1.04
(%)	(0.4)	(6.9)	(26.4)	(40.3)	(20.0)	(4.3)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	75.93	83.53	96.06	103.14	106.43	107.15	105.86	108.19	105.23	131.72	101.91	165.14	101.92		
標準偏差	26.02	27.34	25.13	22.83	21.77	21.26	22.62	20.69	30.64	31.46	23.84	91.82	23.87		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査