

表 2662 体重減少量(kg) %クレアチニン産生速度(%)別 (血液透析濾過患者 女性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
60未満	10	112	282	187	38	3					632		632	1.68	0.84
(%)	(1.6)	(17.7)	(44.6)	(29.6)	(6.0)	(0.5)					(100.0)				
60～	8	54	215	187	49	3	1			1	518		518	1.93	1.08
(%)	(1.5)	(10.4)	(41.5)	(36.1)	(9.5)	(0.6)	(0.2)			(0.2)	(100.0)				
70～	6	84	333	379	88	16	1				907		907	2.02	0.85
(%)	(0.7)	(9.3)	(36.7)	(41.8)	(9.7)	(1.8)	(0.1)				(100.0)				
80～	5	91	468	577	170	23	3		2		1,339		1,339	2.15	0.86
(%)	(0.4)	(6.8)	(35.0)	(43.1)	(12.7)	(1.7)	(0.2)		(0.1)		(100.0)				
90～	6	87	562	925	267	40	5				1,892		1,892	2.24	0.80
(%)	(0.3)	(4.6)	(29.7)	(48.9)	(14.1)	(2.1)	(0.3)				(100.0)				
100～	1	69	667	1,140	327	34	4				2,242		2,242	2.26	0.74
(%)	(0.0)	(3.1)	(29.8)	(50.8)	(14.6)	(1.5)	(0.2)				(100.0)				
110～	1	65	548	1,060	333	35	3	2			2,047		2,047	2.33	0.74
(%)	(0.0)	(3.2)	(26.8)	(51.8)	(16.3)	(1.7)	(0.1)	(0.1)			(100.0)				
120～		33	384	826	245	31	1	1		1	1,522		1,522	2.36	0.76
(%)		(2.2)	(25.2)	(54.3)	(16.1)	(2.0)	(0.1)	(0.1)		(0.1)	(100.0)				
130～	1	14	196	474	132	7	2	3			829		829	2.40	0.72
(%)	(0.1)	(1.7)	(23.6)	(57.2)	(15.9)	(0.8)	(0.2)	(0.4)			(100.0)				
140～		5	116	239	80	16	4	1		6	467	1	468	2.58	1.36
(%)		(1.1)	(24.8)	(51.2)	(17.1)	(3.4)	(0.9)	(0.2)		(1.3)	(100.0)				
合計	38	614	3,771	5,994	1,729	208	24	7	2	8	12,395	1	12,396	2.23	0.84
(%)	(0.3)	(5.0)	(30.4)	(48.4)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
記載なし	1	45	188	210	62	5	1		1		513	286	799	2.06	0.85
(%)	(0.2)	(8.8)	(36.6)	(40.9)	(12.1)	(1.0)	(0.2)		(0.2)		(100.0)				
総計	39	659	3,959	6,204	1,791	213	25	7	3	8	12,908	287	13,195	2.22	0.84
(%)	(0.3)	(5.1)	(30.7)	(48.1)	(13.9)	(1.7)	(0.2)	(0.1)	(0.0)	(0.1)	(100.0)				
平均	72.95	85.10	98.12	105.32	106.73	106.71	114.25	131.03	88.30	144.60	102.30	199.80	102.31		
標準偏差	23.95	26.54	25.09	22.05	21.22	23.07	36.20	14.45	0.28	37.37	23.88		23.89		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査