

表 2666 体重減少量(kg) 透析前リン濃度(mg/dL)別(血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
3未満	6	108	326	360	139	18	3				960	8	968	2.07	0.94
(%)	(0.6)	(11.3)	(34.0)	(37.5)	(14.5)	(1.9)	(0.3)				(100.0)				
3.0～	9	101	420	537	199	33	5	1	1	1	1,307	7	1,314	2.19	1.02
(%)	(0.7)	(7.7)	(32.1)	(41.1)	(15.2)	(2.5)	(0.4)	(0.1)	(0.1)	(0.1)	(100.0)				
3.5～	14	139	689	1,066	417	74	10	1		2	2,412	10	2,422	2.29	1.04
(%)	(0.6)	(5.8)	(28.6)	(44.2)	(17.3)	(3.1)	(0.4)	(0.0)		(0.1)	(100.0)				
4.0～	14	203	1,002	1,757	783	142	29	4		6	3,940	22	3,962	2.39	0.98
(%)	(0.4)	(5.2)	(25.4)	(44.6)	(19.9)	(3.6)	(0.7)	(0.1)		(0.2)	(100.0)				
4.5～	9	195	1,220	2,195	1,076	218	42	6	2	6	4,969	22	4,991	2.47	0.98
(%)	(0.2)	(3.9)	(24.6)	(44.2)	(21.7)	(4.4)	(0.8)	(0.1)	(0.0)	(0.1)	(100.0)				
5.0～	18	211	1,192	2,433	1,256	279	32	7		3	5,431	29	5,460	2.51	0.95
(%)	(0.3)	(3.9)	(21.9)	(44.8)	(23.1)	(5.1)	(0.6)	(0.1)		(0.1)	(100.0)				
5.5～	7	131	918	2,099	1,104	324	49	5		1	4,638	30	4,668	2.61	0.95
(%)	(0.2)	(2.8)	(19.8)	(45.3)	(23.8)	(7.0)	(1.1)	(0.1)		(0.0)	(100.0)				
6.0～		28	175	383	198	46	6	1		2	839	5	844	2.59	1.02
(%)		(3.3)	(20.9)	(45.6)	(23.6)	(5.5)	(0.7)	(0.1)		(0.2)	(100.0)				
6.1～	6	51	495	1,232	762	258	49	3	3	2	2,861	23	2,884	2.75	1.00
(%)	(0.2)	(1.8)	(17.3)	(43.1)	(26.6)	(9.0)	(1.7)	(0.1)	(0.1)	(0.1)	(100.0)				
6.5～	5	54	378	1,020	727	223	42	10	3		2,462	12	2,474	2.81	0.97
(%)	(0.2)	(2.2)	(15.4)	(41.4)	(29.5)	(9.1)	(1.7)	(0.4)	(0.1)		(100.0)				
7.0～	5	28	223	607	499	180	27	4	1	2	1,576	11	1,587	2.90	1.04
(%)	(0.3)	(1.8)	(14.1)	(38.5)	(31.7)	(11.4)	(1.7)	(0.3)	(0.1)	(0.1)	(100.0)				
7.5～		19	114	374	313	122	23	4	2	2	973	5	978	3.00	1.09
(%)		(2.0)	(11.7)	(38.4)	(32.2)	(12.5)	(2.4)	(0.4)	(0.2)	(0.2)	(100.0)				
8.0～	2	10	67	194	189	90	16	4	1	1	574	4	578	3.06	1.11
(%)	(0.3)	(1.7)	(11.7)	(33.8)	(32.9)	(15.7)	(2.8)	(0.7)	(0.2)	(0.2)	(100.0)				
8.5～	1	3	45	117	117	60	16				359	5	364	3.13	1.05
(%)	(0.3)	(0.8)	(12.5)	(32.6)	(32.6)	(16.7)	(4.5)				(100.0)				
9.0～	1	10	56	163	188	114	51	9	3	2	597	4	601	3.41	1.63
(%)	(0.2)	(1.7)	(9.4)	(27.3)	(31.5)	(19.1)	(8.5)	(1.5)	(0.5)	(0.3)	(100.0)				
合計	97	1,291	7,320	14,537	7,967	2,181	400	59	16	30	33,898	197	34,095	2.57	1.03
(%)	(0.3)	(3.8)	(21.6)	(42.9)	(23.5)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし		29	80	117	47	10					283	593	876	2.21	0.94
(%)		(10.2)	(28.3)	(41.3)	(16.6)	(3.5)					(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	4.85	4.73	5.02	5.31	5.64	6.13	6.53	6.65	7.17	5.51	5.37	5.51	5.37		
標準偏差	1.46	1.37	1.31	1.31	1.45	1.58	1.93	2.17	2.12	1.69	1.42	1.50	1.42		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査