

表 2669 体重減少量(kg) 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前ヘモグロビン濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
5未満		1	4	12	4	1	2				24		24	2.67	1.07
(%)		(4.2)	(16.7)	(50.0)	(16.7)	(4.2)	(8.3)				(100.0)				
5～		3	11	9	2	3					28		28	2.18	1.17
(%)		(10.7)	(39.3)	(32.1)	(7.1)	(10.7)					(100.0)				
6～	1	9	25	27	29	2					93	1	94	2.32	0.96
(%)	(1.1)	(9.7)	(26.9)	(29.0)	(31.2)	(2.2)					(100.0)				
7～	3	23	85	153	62	16	8	1	1	2	354	7	361	2.49	1.31
(%)	(0.8)	(6.5)	(24.0)	(43.2)	(17.5)	(4.5)	(2.3)	(0.3)	(0.3)	(0.6)	(100.0)				
8～	5	60	342	552	280	77	21	3		2	1,342	21	1,363	2.48	1.06
(%)	(0.4)	(4.5)	(25.5)	(41.1)	(20.9)	(5.7)	(1.6)	(0.2)		(0.1)	(100.0)				
9～	16	211	1,176	2,185	1,112	281	52	10	2	4	5,049	34	5,083	2.51	1.02
(%)	(0.3)	(4.2)	(23.3)	(43.3)	(22.0)	(5.6)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
10～	34	429	2,443	4,899	2,640	739	116	21	4	12	11,337	52	11,389	2.57	1.05
(%)	(0.3)	(3.8)	(21.5)	(43.2)	(23.3)	(6.5)	(1.0)	(0.2)	(0.0)	(0.1)	(100.0)				
11～	23	349	2,122	4,451	2,445	651	118	16	7	5	10,187	47	10,234	2.59	0.98
(%)	(0.2)	(3.4)	(20.8)	(43.7)	(24.0)	(6.4)	(1.2)	(0.2)	(0.1)	(0.0)	(100.0)				
12～	11	141	760	1,600	974	271	46	6	2	4	3,815	22	3,837	2.62	1.06
(%)	(0.3)	(3.7)	(19.9)	(41.9)	(25.5)	(7.1)	(1.2)	(0.2)	(0.1)	(0.1)	(100.0)				
13～	4	34	199	425	284	86	23	1			1,056	9	1,065	2.68	1.01
(%)	(0.4)	(3.2)	(18.8)	(40.2)	(26.9)	(8.1)	(2.2)	(0.1)			(100.0)				
14～		20	108	182	107	44	11	1		1	474	4	478	2.68	1.17
(%)		(4.2)	(22.8)	(38.4)	(22.6)	(9.3)	(2.3)	(0.2)		(0.2)	(100.0)				
合計	97	1,280	7,275	14,495	7,939	2,171	397	59	16	30	33,759	197	33,956	2.57	1.03
(%)	(0.3)	(3.8)	(21.5)	(42.9)	(23.5)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし		40	125	159	75	20	3				422	593	1,015	2.27	1.01
(%)		(9.5)	(29.6)	(37.7)	(17.8)	(4.7)	(0.7)				(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	10.64	10.75	10.79	10.85	10.92	10.97	10.97	10.72	10.82	10.51	10.85	10.63	10.85		
標準偏差	1.34	1.38	1.29	1.23	1.23	1.26	1.53	1.27	1.20	1.48	1.26	1.51	1.26		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査