

表 2671 体重減少量(kg) HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

HDL-C濃度	0.0未満	0.0～	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	合計	記載なし	総計	平均	標準偏差
20未満	1	7	21	21	15	3					68		68	2.26	1.06
(%)	(1.5)	(10.3)	(30.9)	(30.9)	(22.1)	(4.4)					(100.0)				
20～	4	65	308	612	425	143	32	6	1	1	1,597	4	1,601	2.70	1.10
(%)	(0.3)	(4.1)	(19.3)	(38.3)	(26.6)	(9.0)	(2.0)	(0.4)	(0.1)	(0.1)	(100.0)				
30～	15	193	1,020	2,177	1,275	409	96	10	1	7	5,203	33	5,236	2.65	1.04
(%)	(0.3)	(3.7)	(19.6)	(41.8)	(24.5)	(7.9)	(1.8)	(0.2)	(0.0)	(0.1)	(100.0)				
40～	12	223	1,415	2,732	1,485	434	73	11	3	3	6,391	30	6,421	2.58	1.00
(%)	(0.2)	(3.5)	(22.1)	(42.7)	(23.2)	(6.8)	(1.1)	(0.2)	(0.0)	(0.0)	(100.0)				
50～	15	197	1,163	2,311	1,206	287	41	9	4	6	5,239	18	5,257	2.54	1.03
(%)	(0.3)	(3.8)	(22.2)	(44.1)	(23.0)	(5.5)	(0.8)	(0.2)	(0.1)	(0.1)	(100.0)				
60～	8	145	765	1,557	761	176	24	4	2	4	3,446	12	3,458	2.50	1.04
(%)	(0.2)	(4.2)	(22.2)	(45.2)	(22.1)	(5.1)	(0.7)	(0.1)	(0.1)	(0.1)	(100.0)				
70～	6	70	436	806	381	100	19	2	1	2	1,823	8	1,831	2.50	1.02
(%)	(0.3)	(3.8)	(23.9)	(44.2)	(20.9)	(5.5)	(1.0)	(0.1)	(0.1)	(0.1)	(100.0)				
80～	7	43	217	401	210	45	3	1			927	1	928	2.45	0.93
(%)	(0.8)	(4.6)	(23.4)	(43.3)	(22.7)	(4.9)	(0.3)	(0.1)			(100.0)				
90～	3	24	78	173	90	21	4	1			394	1	395	2.48	1.02
(%)	(0.8)	(6.1)	(19.8)	(43.9)	(22.8)	(5.3)	(1.0)	(0.3)			(100.0)				
100～		16	77	149	76	18	6	1			343	1	344	2.51	0.99
(%)		(4.7)	(22.4)	(43.4)	(22.2)	(5.2)	(1.7)	(0.3)			(100.0)				
合計	71	983	5,500	10,939	5,924	1,636	298	45	12	23	25,431	108	25,539	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(43.0)	(23.3)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
記載なし	26	337	1,900	3,715	2,090	555	102	14	4	7	8,750	682	9,432	2.57	1.04
(%)	(0.3)	(3.9)	(21.7)	(42.5)	(23.9)	(6.3)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
総計	97	1,320	7,400	14,654	8,014	2,191	400	59	16	30	34,181	790	34,971	2.57	1.03
(%)	(0.3)	(3.9)	(21.6)	(42.9)	(23.4)	(6.4)	(1.2)	(0.2)	(0.0)	(0.1)	(100.0)				
平均	53.28	52.12	51.88	51.73	50.52	48.31	46.71	47.87	49.75	48.91	51.21	48.49	51.20		
標準偏差	19.35	18.16	17.22	17.15	17.37	16.89	18.83	18.43	13.30	12.78	17.30	16.03	17.29		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査