

表 2685 体重減少率(%) Body Mass Index別 (施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上)

Body Mass Index	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
14未満	21	18	42	47	56	52	42	23	8	5	314	28	342	4.51	2.27
(%)	(6.7)	(5.7)	(13.4)	(15.0)	(17.8)	(16.6)	(13.4)	(7.3)	(2.5)	(1.6)	(100.0)				
14～	135	118	257	370	492	448	347	186	60	51	2,464	7	2,471	4.67	2.14
(%)	(5.5)	(4.8)	(10.4)	(15.0)	(20.0)	(18.2)	(14.1)	(7.5)	(2.4)	(2.1)	(100.0)				
16～	270	419	857	1,550	2,250	2,240	1,458	668	252	125	10,089	28	10,117	4.81	1.85
(%)	(2.7)	(4.2)	(8.5)	(15.4)	(22.3)	(22.2)	(14.5)	(6.6)	(2.5)	(1.2)	(100.0)				
18～	503	759	1,862	3,633	5,020	4,762	2,702	1,081	334	176	20,832	32	20,864	4.69	1.72
(%)	(2.4)	(3.6)	(8.9)	(17.4)	(24.1)	(22.9)	(13.0)	(5.2)	(1.6)	(0.8)	(100.0)				
20～	502	960	2,358	4,965	6,855	5,631	2,772	934	262	150	25,389	19	25,408	4.54	1.61
(%)	(2.0)	(3.8)	(9.3)	(19.6)	(27.0)	(22.2)	(10.9)	(3.7)	(1.0)	(0.6)	(100.0)				
22～	385	737	2,080	4,363	5,973	4,170	1,757	535	116	67	20,183	7	20,190	4.39	1.49
(%)	(1.9)	(3.7)	(10.3)	(21.6)	(29.6)	(20.7)	(8.7)	(2.7)	(0.6)	(0.3)	(100.0)				
24～	190	436	1,280	2,854	3,694	2,320	873	226	58	29	11,960	4	11,964	4.30	1.42
(%)	(1.6)	(3.6)	(10.7)	(23.9)	(30.9)	(19.4)	(7.3)	(1.9)	(0.5)	(0.2)	(100.0)				
26～	94	228	647	1,582	1,937	1,177	420	96	23	7	6,211	3	6,214	4.26	1.35
(%)	(1.5)	(3.7)	(10.4)	(25.5)	(31.2)	(19.0)	(6.8)	(1.5)	(0.4)	(0.1)	(100.0)				
28～	48	89	316	780	1,012	563	187	41	10	1	3,047	3	3,050	4.24	1.30
(%)	(1.6)	(2.9)	(10.4)	(25.6)	(33.2)	(18.5)	(6.1)	(1.3)	(0.3)	(0.0)	(100.0)				
30～	20	69	220	608	818	451	149	45	8	6	2,394	4	2,398	4.35	1.32
(%)	(0.8)	(2.9)	(9.2)	(25.4)	(34.2)	(18.8)	(6.2)	(1.9)	(0.3)	(0.3)	(100.0)				
合計	2,168	3,833	9,919	20,752	28,107	21,814	10,707	3,835	1,131	617	102,883	135	103,018	4.51	1.61
(%)	(2.1)	(3.7)	(9.6)	(20.2)	(27.3)	(21.2)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
記載なし	184	335	712	1,369	1,750	1,308	633	250	65	47	6,653	2,844	9,497	4.39	1.69
(%)	(2.8)	(5.0)	(10.7)	(20.6)	(26.3)	(19.7)	(9.5)	(3.8)	(1.0)	(0.7)	(100.0)				
総計	2,352	4,168	10,631	22,121	29,857	23,122	11,340	4,085	1,196	664	109,536	2,979	112,515	4.50	1.62
(%)	(2.1)	(3.8)	(9.7)	(20.2)	(27.3)	(21.1)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
平均	20.84	21.51	21.92	22.27	22.17	21.57	20.91	20.28	19.86	19.63	21.75	17.71	21.74		
標準偏差	3.41	3.43	3.49	3.54	3.52	3.36	3.22	3.14	2.97	2.93	3.48	5.78	3.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査