

表 2686 体重減少率(%) Body Mass Index別 (施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上)

Body Mass Index	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
14未満	43	40	61	93	143	116	115	69	31	22	733	21	754	4.91	2.40
(%)	(5.9)	(5.5)	(8.3)	(12.7)	(19.5)	(15.8)	(15.7)	(9.4)	(4.2)	(3.0)	(100.0)				
14～	127	135	301	496	714	813	627	373	168	97	3,851	7	3,858	5.09	2.05
(%)	(3.3)	(3.5)	(7.8)	(12.9)	(18.5)	(21.1)	(16.3)	(9.7)	(4.4)	(2.5)	(100.0)				
16～	227	354	767	1,476	2,138	2,168	1,555	799	290	152	9,926	6	9,932	4.97	1.87
(%)	(2.3)	(3.6)	(7.7)	(14.9)	(21.5)	(21.8)	(15.7)	(8.0)	(2.9)	(1.5)	(100.0)				
18～	244	495	1,188	2,296	3,482	3,202	1,905	765	241	127	13,945	7	13,952	4.77	1.70
(%)	(1.7)	(3.5)	(8.5)	(16.5)	(25.0)	(23.0)	(13.7)	(5.5)	(1.7)	(0.9)	(100.0)				
20～	249	460	1,152	2,374	3,245	2,734	1,266	482	126	66	12,154	8	12,162	4.54	1.62
(%)	(2.0)	(3.8)	(9.5)	(19.5)	(26.7)	(22.5)	(10.4)	(4.0)	(1.0)	(0.5)	(100.0)				
22～	149	310	916	1,887	2,470	1,673	661	210	49	23	8,348	4	8,352	4.34	1.49
(%)	(1.8)	(3.7)	(11.0)	(22.6)	(29.6)	(20.0)	(7.9)	(2.5)	(0.6)	(0.3)	(100.0)				
24～	85	216	612	1,293	1,468	896	282	78	13	9	4,952	4	4,956	4.15	1.40
(%)	(1.7)	(4.4)	(12.4)	(26.1)	(29.6)	(18.1)	(5.7)	(1.6)	(0.3)	(0.2)	(100.0)				
26～	58	101	350	777	897	433	131	29	8	3	2,787	4	2,791	4.08	1.34
(%)	(2.1)	(3.6)	(12.6)	(27.9)	(32.2)	(15.5)	(4.7)	(1.0)	(0.3)	(0.1)	(100.0)				
28～	28	73	221	466	486	170	68	5	4	3	1,524		1,524	3.92	1.33
(%)	(1.8)	(4.8)	(14.5)	(30.6)	(31.9)	(11.2)	(4.5)	(0.3)	(0.3)	(0.2)	(100.0)				
30～	22	57	218	488	437	188	47	5		2	1,464	2	1,466	3.91	1.22
(%)	(1.5)	(3.9)	(14.9)	(33.3)	(29.8)	(12.8)	(3.2)	(0.3)		(0.1)	(100.0)				
合計	1,232	2,241	5,786	11,646	15,480	12,393	6,657	2,815	930	504	59,684	63	59,747	4.59	1.70
(%)	(2.1)	(3.8)	(9.7)	(19.5)	(25.9)	(20.8)	(11.2)	(4.7)	(1.6)	(0.8)	(100.0)				
記載なし	112	200	504	817	1,078	774	430	208	69	37	4,229	1,718	5,947	4.46	1.78
(%)	(2.6)	(4.7)	(11.9)	(19.3)	(25.5)	(18.3)	(10.2)	(4.9)	(1.6)	(0.9)	(100.0)				
総計	1,344	2,441	6,290	12,463	16,558	13,167	7,087	3,023	999	541	63,913	1,781	65,694	4.58	1.70
(%)	(2.1)	(3.8)	(9.8)	(19.5)	(25.9)	(20.6)	(11.1)	(4.7)	(1.6)	(0.8)	(100.0)				
平均	20.09	20.88	21.48	21.73	21.26	20.35	19.43	18.69	18.12	18.02	20.75	16.94	20.74		
標準偏差	3.97	3.92	4.06	4.07	3.84	3.42	3.15	2.79	2.69	2.94	3.82	6.83	3.82		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査