

表 2687 体重減少率(%) 透析前BUN(mg/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前BUN	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	94	84	81	89	67	31	13	4	4	1	468	11	479	2.73	1.97
(%)	(20.1)	(17.9)	(17.3)	(19.0)	(14.3)	(6.6)	(2.8)	(0.9)	(0.9)	(0.2)	(100.0)				
20～	302	330	521	494	389	255	103	30	15	7	2,446	34	2,480	3.20	1.87
(%)	(12.3)	(13.5)	(21.3)	(20.2)	(15.9)	(10.4)	(4.2)	(1.2)	(0.6)	(0.3)	(100.0)				
30～	590	939	1,738	2,391	2,205	1,345	671	253	77	60	10,269	89	10,358	3.81	1.80
(%)	(5.7)	(9.1)	(16.9)	(23.3)	(21.5)	(13.1)	(6.5)	(2.5)	(0.7)	(0.6)	(100.0)				
40～	862	1,522	3,599	6,087	6,728	4,561	2,094	779	266	143	26,641	170	26,811	4.18	1.68
(%)	(3.2)	(5.7)	(13.5)	(22.8)	(25.3)	(17.1)	(7.9)	(2.9)	(1.0)	(0.5)	(100.0)				
50～	786	1,606	4,448	9,155	11,579	8,617	3,998	1,482	446	256	42,373	286	42,659	4.45	1.60
(%)	(1.9)	(3.8)	(10.5)	(21.6)	(27.3)	(20.3)	(9.4)	(3.5)	(1.1)	(0.6)	(100.0)				
60～	558	1,189	3,519	8,450	12,131	9,289	4,761	1,710	494	286	42,387	256	42,643	4.64	1.55
(%)	(1.3)	(2.8)	(8.3)	(19.9)	(28.6)	(21.9)	(11.2)	(4.0)	(1.2)	(0.7)	(100.0)				
70～	269	568	1,881	4,901	7,962	6,866	3,654	1,417	426	217	28,161	161	28,322	4.85	1.53
(%)	(1.0)	(2.0)	(6.7)	(17.4)	(28.3)	(24.4)	(13.0)	(5.0)	(1.5)	(0.8)	(100.0)				
80～	120	191	675	1,962	3,575	3,405	1,946	854	263	134	13,125	87	13,212	5.05	1.55
(%)	(0.9)	(1.5)	(5.1)	(14.9)	(27.2)	(25.9)	(14.8)	(6.5)	(2.0)	(1.0)	(100.0)				
90～	50	70	244	635	1,187	1,264	774	370	120	61	4,775	28	4,803	5.16	1.60
(%)	(1.0)	(1.5)	(5.1)	(13.3)	(24.9)	(26.5)	(16.2)	(7.7)	(2.5)	(1.3)	(100.0)				
100～	12	22	74	149	276	371	236	122	48	25	1,335	13	1,348	5.35	1.67
(%)	(0.9)	(1.6)	(5.5)	(11.2)	(20.7)	(27.8)	(17.7)	(9.1)	(3.6)	(1.9)	(100.0)				
110～	6	10	21	46	75	92	78	36	21	8	393	1	394	5.35	1.82
(%)	(1.5)	(2.5)	(5.3)	(11.7)	(19.1)	(23.4)	(19.8)	(9.2)	(5.3)	(2.0)	(100.0)				
120～	2	11	10	29	35	41	40	23	5	5	201	3	204	5.26	1.89
(%)	(1.0)	(5.5)	(5.0)	(14.4)	(17.4)	(20.4)	(19.9)	(11.4)	(2.5)	(2.5)	(100.0)				
合計	3,651	6,542	16,811	34,388	46,209	36,137	18,368	7,080	2,185	1,203	172,574	1,139	173,713	4.54	1.65
(%)	(2.1)	(3.8)	(9.7)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
記載なし	45	67	110	196	206	152	59	28	10	2	875	3,621	4,496	4.05	1.77
(%)	(5.1)	(7.7)	(12.6)	(22.4)	(23.5)	(17.4)	(6.7)	(3.2)	(1.1)	(0.2)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	50.18	52.48	55.63	58.92	61.67	63.60	64.92	66.01	66.16	65.23	60.95	59.29	60.94		
標準偏差	17.02	16.27	15.34	14.80	14.67	15.12	15.60	16.27	17.18	17.01	15.63	16.90	15.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査