

表 2688 体重減少率(%) 透析前クレアチニン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前クレアチニン濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
2未満	13	5	15	17	10	11	2		2	1	76		76	3.33	2.16
(%)	(17.1)	(6.6)	(19.7)	(22.4)	(13.2)	(14.5)	(2.6)		(2.6)	(1.3)	(100.0)				
2～	196	203	279	345	290	166	104	33	8	8	1,632	24	1,656	3.40	1.99
(%)	(12.0)	(12.4)	(17.1)	(21.1)	(17.8)	(10.2)	(6.4)	(2.0)	(0.5)	(0.5)	(100.0)				
4～	590	749	1,215	1,565	1,490	1,005	580	284	130	96	7,704	91	7,795	3.88	2.06
(%)	(7.7)	(9.7)	(15.8)	(20.3)	(19.3)	(13.0)	(7.5)	(3.7)	(1.7)	(1.2)	(100.0)				
6～	942	1,416	2,895	4,615	5,376	4,207	2,471	1,162	448	280	23,812	194	24,006	4.39	1.91
(%)	(4.0)	(5.9)	(12.2)	(19.4)	(22.6)	(17.7)	(10.4)	(4.9)	(1.9)	(1.2)	(100.0)				
8～	883	1,835	4,664	9,022	11,863	9,442	5,069	2,089	624	371	45,862	253	46,115	4.56	1.68
(%)	(1.9)	(4.0)	(10.2)	(19.7)	(25.9)	(20.6)	(11.1)	(4.6)	(1.4)	(0.8)	(100.0)				
10～	554	1,358	4,287	9,769	13,422	10,513	5,157	1,901	536	245	47,742	292	48,034	4.61	1.52
(%)	(1.2)	(2.8)	(9.0)	(20.5)	(28.1)	(22.0)	(10.8)	(4.0)	(1.1)	(0.5)	(100.0)				
12～	299	672	2,318	5,979	8,773	6,857	3,203	1,043	286	141	29,571	183	29,754	4.65	1.46
(%)	(1.0)	(2.3)	(7.8)	(20.2)	(29.7)	(23.2)	(10.8)	(3.5)	(1.0)	(0.5)	(100.0)				
14～	126	223	866	2,294	3,690	2,907	1,300	416	109	38	11,969	67	12,036	4.68	1.42
(%)	(1.1)	(1.9)	(7.2)	(19.2)	(30.8)	(24.3)	(10.9)	(3.5)	(0.9)	(0.3)	(100.0)				
16～	45	78	262	764	1,259	992	464	145	38	19	4,066	34	4,100	4.72	1.45
(%)	(1.1)	(1.9)	(6.4)	(18.8)	(31.0)	(24.4)	(11.4)	(3.6)	(0.9)	(0.5)	(100.0)				
合計	3,648	6,539	16,801	34,370	46,173	36,100	18,350	7,073	2,181	1,199	172,434	1,138	173,572	4.54	1.65
(%)	(2.1)	(3.8)	(9.7)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
記載なし	48	70	120	214	242	189	77	35	14	6	1,015	3,622	4,637	4.19	1.79
(%)	(4.7)	(6.9)	(11.8)	(21.1)	(23.8)	(18.6)	(7.6)	(3.4)	(1.4)	(0.6)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	8.41	9.01	9.70	10.29	10.60	10.64	10.42	10.10	9.74	9.38	10.29	9.95	10.29		
標準偏差	3.12	2.92	2.80	2.75	2.72	2.71	2.68	2.69	2.67	2.74	2.78	3.15	2.79		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査