

表 2690 体重減少率(%) single pool Kt/V(Kt/Vsp)別(施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上)

single pool Kt/V (Kt/Vsp)	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
0.4未満	7	10	13	15	17	7	9	1			79		79	3.56	1.88
(%)	(8.9)	(12.7)	(16.5)	(19.0)	(21.5)	(8.9)	(11.4)	(1.3)			(100.0)				
0.4～	14	19	17	18	27	17	2	3			117		117	3.35	1.81
(%)	(12.0)	(16.2)	(14.5)	(15.4)	(23.1)	(14.5)	(1.7)	(2.6)			(100.0)				
0.6～	100	86	142	111	88	49	14	4	1	1	596		596	2.86	1.77
(%)	(16.8)	(14.4)	(23.8)	(18.6)	(14.8)	(8.2)	(2.3)	(0.7)	(0.2)	(0.2)	(100.0)				
0.8～	322	344	677	947	805	388	136	40	11	2	3,672		3,672	3.44	1.66
(%)	(8.8)	(9.4)	(18.4)	(25.8)	(21.9)	(10.6)	(3.7)	(1.1)	(0.3)	(0.1)	(100.0)				
1.0～	591	937	2,080	3,786	4,237	2,290	851	248	68	30	15,118		15,118	3.96	1.55
(%)	(3.9)	(6.2)	(13.8)	(25.0)	(28.0)	(15.1)	(5.6)	(1.6)	(0.4)	(0.2)	(100.0)				
1.2～	616	1,262	3,411	7,230	9,710	6,807	2,750	850	202	85	32,923	1	32,924	4.37	1.48
(%)	(1.9)	(3.8)	(10.4)	(22.0)	(29.5)	(20.7)	(8.4)	(2.6)	(0.6)	(0.3)	(100.0)				
1.4～	327	783	2,325	5,814	8,546	7,315	3,748	1,249	350	164	30,621	2	30,623	4.72	1.50
(%)	(1.1)	(2.6)	(7.6)	(19.0)	(27.9)	(23.9)	(12.2)	(4.1)	(1.1)	(0.5)	(100.0)				
1.6～	106	284	865	2,174	3,586	3,537	2,114	879	268	157	13,970	5	13,975	5.01	1.58
(%)	(0.8)	(2.0)	(6.2)	(15.6)	(25.7)	(25.3)	(15.1)	(6.3)	(1.9)	(1.1)	(100.0)				
1.8～	46	71	244	604	989	1,125	746	375	135	108	4,443	11	4,454	5.28	1.81
(%)	(1.0)	(1.6)	(5.5)	(13.6)	(22.3)	(25.3)	(16.8)	(8.4)	(3.0)	(2.4)	(100.0)				
2.0～	15	37	113	203	412	417	344	220	92	80	1,933	47	1,980	5.55	2.04
(%)	(0.8)	(1.9)	(5.8)	(10.5)	(21.3)	(21.6)	(17.8)	(11.4)	(4.8)	(4.1)	(100.0)				
合計	2,144	3,833	9,887	20,902	28,417	21,952	10,714	3,869	1,127	627	103,472	66	103,538	4.52	1.61
(%)	(2.1)	(3.7)	(9.6)	(20.2)	(27.5)	(21.2)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
記載なし	208	335	744	1,219	1,440	1,170	626	216	69	37	6,064	2,913	8,977	4.32	1.74
(%)	(3.4)	(5.5)	(12.3)	(20.1)	(23.7)	(19.3)	(10.3)	(3.6)	(1.1)	(0.6)	(100.0)				
総計	2,352	4,168	10,631	22,121	29,857	23,122	11,340	4,085	1,196	664	109,536	2,979	112,515	4.50	1.62
(%)	(2.1)	(3.8)	(9.7)	(20.2)	(27.3)	(21.1)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
平均	1.22	1.28	1.32	1.36	1.39	1.44	1.49	1.54	1.58	1.66	1.40	2.20	1.40		
標準偏差	0.28	0.26	0.25	0.24	0.24	0.24	0.25	0.27	0.28	0.31	0.25	0.38	0.26		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査