

表 2691 体重減少率(%) single pool Kt/V(Kt/Vsp)別(施設血液透析患者 女性のみ 透析回数週3回 透析歴2年以上)

single pool Kt/V (Kt/Vsp)	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
0.4未満	1	4	2	2	6	2	4				21		21	3.95	1.91
(%)	(4.8)	(19.0)	(9.5)	(9.5)	(28.6)	(9.5)	(19.0)				(100.0)				
0.4～	9	4	10	13	10	2	1	1			50		50	3.03	1.64
(%)	(18.0)	(8.0)	(20.0)	(26.0)	(20.0)	(4.0)	(2.0)	(2.0)			(100.0)				
0.6～	22	26	35	34	34	15	10	2			178		178	3.15	1.85
(%)	(12.4)	(14.6)	(19.7)	(19.1)	(19.1)	(8.4)	(5.6)	(1.1)			(100.0)				
0.8～	78	83	174	205	153	77	45	11	5	1	832		832	3.40	1.78
(%)	(9.4)	(10.0)	(20.9)	(24.6)	(18.4)	(9.3)	(5.4)	(1.3)	(0.6)	(0.1)	(100.0)				
1.0～	201	276	534	861	733	381	158	48	14	4	3,210		3,210	3.66	1.67
(%)	(6.3)	(8.6)	(16.6)	(26.8)	(22.8)	(11.9)	(4.9)	(1.5)	(0.4)	(0.1)	(100.0)				
1.2～	297	481	1,251	2,083	2,332	1,382	611	215	42	33	8,727		8,727	4.07	1.61
(%)	(3.4)	(5.5)	(14.3)	(23.9)	(26.7)	(15.8)	(7.0)	(2.5)	(0.5)	(0.4)	(100.0)				
1.4～	274	626	1,665	3,270	4,314	3,132	1,515	505	154	58	15,513		15,513	4.43	1.56
(%)	(1.8)	(4.0)	(10.7)	(21.1)	(27.8)	(20.2)	(9.8)	(3.3)	(1.0)	(0.4)	(100.0)				
1.6～	189	427	1,214	2,830	4,240	3,629	1,931	831	252	117	15,660		15,660	4.77	1.59
(%)	(1.2)	(2.7)	(7.8)	(18.1)	(27.1)	(23.2)	(12.3)	(5.3)	(1.6)	(0.7)	(100.0)				
1.8～	102	194	669	1,538	2,423	2,362	1,423	671	229	123	9,734	3	9,737	5.02	1.68
(%)	(1.0)	(2.0)	(6.9)	(15.8)	(24.9)	(24.3)	(14.6)	(6.9)	(2.4)	(1.3)	(100.0)				
2.0～	50	108	299	879	1,485	1,528	1,042	578	240	158	6,367	19	6,386	5.32	1.76
(%)	(0.8)	(1.7)	(4.7)	(13.8)	(23.3)	(24.0)	(16.4)	(9.1)	(3.8)	(2.5)	(100.0)				
合計	1,223	2,229	5,853	11,715	15,730	12,510	6,740	2,862	936	494	60,292	22	60,314	4.59	1.69
(%)	(2.0)	(3.7)	(9.7)	(19.4)	(26.1)	(20.7)	(11.2)	(4.7)	(1.6)	(0.8)	(100.0)				
記載なし	121	212	437	748	828	657	347	161	63	47	3,621	1,759	5,380	4.39	1.85
(%)	(3.3)	(5.9)	(12.1)	(20.7)	(22.9)	(18.1)	(9.6)	(4.4)	(1.7)	(1.3)	(100.0)				
総計	1,344	2,441	6,290	12,463	16,558	13,167	7,087	3,023	999	541	63,913	1,781	65,694	4.58	1.70
(%)	(2.1)	(3.8)	(9.8)	(19.5)	(25.9)	(20.6)	(11.1)	(4.7)	(1.6)	(0.8)	(100.0)				
平均	1.41	1.47	1.51	1.57	1.62	1.67	1.71	1.77	1.82	1.87	1.62	2.36	1.62		
標準偏差	0.33	0.32	0.30	0.30	0.29	0.29	0.30	0.31	0.32	0.33	0.31	0.34	0.31		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査