

表 2693 体重減少率(%) equilibrated Kt/V(eKt/V)別(施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上)

equilibrated Kt/V (eKt/V)	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
0.4未満	9	13	15	20	23	10	10	3			103		103	3.62	1.90
(%)	(8.7)	(12.6)	(14.6)	(19.4)	(22.3)	(9.7)	(9.7)	(2.9)			(100.0)				
0.4～	41	44	52	33	46	28	8	5	1		258	1	259	3.06	1.90
(%)	(15.9)	(17.1)	(20.2)	(12.8)	(17.8)	(10.9)	(3.1)	(1.9)	(0.4)		(100.0)				
0.6～	250	249	468	560	425	202	62	18	6	1	2,241		2,241	3.21	1.66
(%)	(11.2)	(11.1)	(20.9)	(25.0)	(19.0)	(9.0)	(2.8)	(0.8)	(0.3)	(0.0)	(100.0)				
0.8～	555	829	1,785	3,087	3,256	1,685	599	191	43	24	12,054		12,054	3.85	1.57
(%)	(4.6)	(6.9)	(14.8)	(25.6)	(27.0)	(14.0)	(5.0)	(1.6)	(0.4)	(0.2)	(100.0)				
1.0～	685	1,319	3,544	7,356	9,746	6,744	2,709	829	203	90	33,225		33,225	4.34	1.49
(%)	(2.1)	(4.0)	(10.7)	(22.1)	(29.3)	(20.3)	(8.2)	(2.5)	(0.6)	(0.3)	(100.0)				
1.2～	404	945	2,683	6,694	9,808	8,266	4,228	1,415	399	188	35,030	4	35,034	4.70	1.51
(%)	(1.2)	(2.7)	(7.7)	(19.1)	(28.0)	(23.6)	(12.1)	(4.0)	(1.1)	(0.5)	(100.0)				
1.4～	139	327	965	2,356	3,790	3,633	2,152	882	276	159	14,679	10	14,689	4.96	1.61
(%)	(0.9)	(2.2)	(6.6)	(16.1)	(25.8)	(24.7)	(14.7)	(6.0)	(1.9)	(1.1)	(100.0)				
1.6～	38	64	258	591	930	1,024	665	358	125	102	4,155	20	4,175	5.25	1.81
(%)	(0.9)	(1.5)	(6.2)	(14.2)	(22.4)	(24.6)	(16.0)	(8.6)	(3.0)	(2.5)	(100.0)				
1.8～	14	32	81	132	260	240	170	109	54	39	1,131	13	1,144	5.30	2.03
(%)	(1.2)	(2.8)	(7.2)	(11.7)	(23.0)	(21.2)	(15.0)	(9.6)	(4.8)	(3.4)	(100.0)				
2.0～	9	11	35	77	138	121	114	60	20	26	611	36	647	5.44	2.14
(%)	(1.5)	(1.8)	(5.7)	(12.6)	(22.6)	(19.8)	(18.7)	(9.8)	(3.3)	(4.3)	(100.0)				
合計	2,144	3,833	9,886	20,906	28,422	21,953	10,717	3,870	1,127	629	103,487	84	103,571	4.52	1.61
(%)	(2.1)	(3.7)	(9.6)	(20.2)	(27.5)	(21.2)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
記載なし	208	335	745	1,215	1,435	1,169	623	215	69	35	6,049	2,895	8,944	4.32	1.74
(%)	(3.4)	(5.5)	(12.3)	(20.1)	(23.7)	(19.3)	(10.3)	(3.6)	(1.1)	(0.6)	(100.0)				
総計	2,352	4,168	10,631	22,121	29,857	23,122	11,340	4,085	1,196	664	109,536	2,979	112,515	4.50	1.62
(%)	(2.1)	(3.8)	(9.7)	(20.2)	(27.3)	(21.1)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
平均	1.06	1.12	1.15	1.19	1.22	1.26	1.30	1.34	1.37	1.44	1.22	2.00	1.23		
標準偏差	0.26	0.25	0.24	0.22	0.22	0.22	0.23	0.25	0.26	0.30	0.23	0.50	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査