

表 2699 体重減少率(%) %クレアチニン産生速度(%)別(施設血液透析患者 男性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	764	810	1,310	1,581	1,456	851	398	171	50	32	7,423	1	7,424	3.51	1.89
(%)	(10.3)	(10.9)	(17.6)	(21.3)	(19.6)	(11.5)	(5.4)	(2.3)	(0.7)	(0.4)	(100.0)				
60～	299	441	809	1,273	1,296	909	473	190	64	31	5,785	1	5,786	4.04	1.81
(%)	(5.2)	(7.6)	(14.0)	(22.0)	(22.4)	(15.7)	(8.2)	(3.3)	(1.1)	(0.5)	(100.0)				
70～	272	491	1,138	1,888	2,198	1,607	812	336	97	69	8,908	1	8,909	4.30	1.74
(%)	(3.1)	(5.5)	(12.8)	(21.2)	(24.7)	(18.0)	(9.1)	(3.8)	(1.1)	(0.8)	(100.0)				
80～	228	544	1,367	2,581	3,299	2,562	1,251	467	153	106	12,558	1	12,559	4.48	1.64
(%)	(1.8)	(4.3)	(10.9)	(20.6)	(26.3)	(20.4)	(10.0)	(3.7)	(1.2)	(0.8)	(100.0)				
90～	212	508	1,479	3,331	4,416	3,349	1,724	663	200	101	15,983	6	15,989	4.59	1.56
(%)	(1.3)	(3.2)	(9.3)	(20.8)	(27.6)	(21.0)	(10.8)	(4.1)	(1.3)	(0.6)	(100.0)				
100～	139	414	1,341	3,403	4,859	3,797	1,887	688	200	88	16,816	2	16,818	4.68	1.49
(%)	(0.8)	(2.5)	(8.0)	(20.2)	(28.9)	(22.6)	(11.2)	(4.1)	(1.2)	(0.5)	(100.0)				
110～	119	301	1,082	2,931	4,420	3,647	1,714	584	173	88	15,059	4	15,063	4.73	1.45
(%)	(0.8)	(2.0)	(7.2)	(19.5)	(29.4)	(24.2)	(11.4)	(3.9)	(1.1)	(0.6)	(100.0)				
120～	57	168	728	2,029	3,292	2,687	1,278	421	102	51	10,813	6	10,819	4.77	1.39
(%)	(0.5)	(1.6)	(6.7)	(18.8)	(30.4)	(24.8)	(11.8)	(3.9)	(0.9)	(0.5)	(100.0)				
130～	29	91	380	1,128	1,910	1,523	723	228	55	25	6,092	7	6,099	4.79	1.37
(%)	(0.5)	(1.5)	(6.2)	(18.5)	(31.4)	(25.0)	(11.9)	(3.7)	(0.9)	(0.4)	(100.0)				
140～	26	64	258	763	1,287	1,032	467	128	34	36	4,095	56	4,151	4.79	1.56
(%)	(0.6)	(1.6)	(6.3)	(18.6)	(31.4)	(25.2)	(11.4)	(3.1)	(0.8)	(0.9)	(100.0)				
合計	2,145	3,832	9,892	20,908	28,433	21,964	10,727	3,876	1,128	627	103,532	85	103,617	4.52	1.61
(%)	(2.1)	(3.7)	(9.6)	(20.2)	(27.5)	(21.2)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
記載なし	207	336	739	1,213	1,424	1,158	613	209	68	37	6,004	2,894	8,898	4.31	1.74
(%)	(3.4)	(5.6)	(12.3)	(20.2)	(23.7)	(19.3)	(10.2)	(3.5)	(1.1)	(0.6)	(100.0)				
総計	2,352	4,168	10,631	22,121	29,857	23,122	11,340	4,085	1,196	664	109,536	2,979	112,515	4.50	1.62
(%)	(2.1)	(3.8)	(9.7)	(20.2)	(27.3)	(21.1)	(10.4)	(3.7)	(1.1)	(0.6)	(100.0)				
平均	71.99	82.24	90.82	97.80	101.84	103.27	102.82	100.72	99.29	98.80	98.94	167.06	99.00		
標準偏差	29.24	28.25	27.10	25.55	24.38	23.37	23.08	22.67	22.32	25.81	25.54	52.48	25.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査