

表 2704 体重減少率(%) 透析前リン濃度(mg/dL)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前リン濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
3未満	463	643	1,054	1,419	1,418	985	518	209	73	41	6,823	66	6,889	3.87	1.90
(%)	(6.8)	(9.4)	(15.4)	(20.8)	(20.8)	(14.4)	(7.6)	(3.1)	(1.1)	(0.6)	(100.0)				
3.0～	370	561	1,093	1,758	2,009	1,290	689	264	78	57	8,169	67	8,236	4.12	1.78
(%)	(4.5)	(6.9)	(13.4)	(21.5)	(24.6)	(15.8)	(8.4)	(3.2)	(1.0)	(0.7)	(100.0)				
3.5～	450	828	1,751	3,079	3,578	2,547	1,167	455	142	89	14,086	108	14,194	4.24	1.71
(%)	(3.2)	(5.9)	(12.4)	(21.9)	(25.4)	(18.1)	(8.3)	(3.2)	(1.0)	(0.6)	(100.0)				
4.0～	535	1,019	2,500	4,576	5,601	4,112	1,957	684	221	107	21,312	140	21,452	4.34	1.64
(%)	(2.5)	(4.8)	(11.7)	(21.5)	(26.3)	(19.3)	(9.2)	(3.2)	(1.0)	(0.5)	(100.0)				
4.5～	507	958	2,689	5,493	7,093	5,263	2,548	939	264	138	25,892	186	26,078	4.47	1.60
(%)	(2.0)	(3.7)	(10.4)	(21.2)	(27.4)	(20.3)	(9.8)	(3.6)	(1.0)	(0.5)	(100.0)				
5.0～	445	864	2,530	5,462	7,375	5,580	2,735	966	311	146	26,414	210	26,624	4.54	1.58
(%)	(1.7)	(3.3)	(9.6)	(20.7)	(27.9)	(21.1)	(10.4)	(3.7)	(1.2)	(0.6)	(100.0)				
5.5～	301	608	1,968	4,574	6,495	5,113	2,578	940	245	152	22,974	148	23,122	4.64	1.55
(%)	(1.3)	(2.6)	(8.6)	(19.9)	(28.3)	(22.3)	(11.2)	(4.1)	(1.1)	(0.7)	(100.0)				
6.0～	56	82	316	759	1,094	913	448	179	60	36	3,943	23	3,966	4.73	1.58
(%)	(1.4)	(2.1)	(8.0)	(19.2)	(27.7)	(23.2)	(11.4)	(4.5)	(1.5)	(0.9)	(100.0)				
6.1～	157	318	1,002	2,490	3,689	3,049	1,518	605	177	87	13,092	73	13,165	4.72	1.54
(%)	(1.2)	(2.4)	(7.7)	(19.0)	(28.2)	(23.3)	(11.6)	(4.6)	(1.4)	(0.7)	(100.0)				
6.5～	113	249	792	1,973	3,183	2,674	1,432	605	184	90	11,295	67	11,362	4.83	1.55
(%)	(1.0)	(2.2)	(7.0)	(17.5)	(28.2)	(23.7)	(12.7)	(5.4)	(1.6)	(0.8)	(100.0)				
7.0～	77	143	439	1,161	1,861	1,751	1,005	392	124	75	7,028	52	7,080	4.93	1.59
(%)	(1.1)	(2.0)	(6.2)	(16.5)	(26.5)	(24.9)	(14.3)	(5.6)	(1.8)	(1.1)	(100.0)				
7.5～	51	78	232	637	1,060	1,058	605	293	85	65	4,164	34	4,198	5.04	1.65
(%)	(1.2)	(1.9)	(5.6)	(15.3)	(25.5)	(25.4)	(14.5)	(7.0)	(2.0)	(1.6)	(100.0)				
8.0～	27	53	140	330	607	619	402	189	70	34	2,471	14	2,485	5.14	1.69
(%)	(1.1)	(2.1)	(5.7)	(13.4)	(24.6)	(25.1)	(16.3)	(7.6)	(2.8)	(1.4)	(100.0)				
8.5～	21	32	86	237	384	385	257	111	49	30	1,592	9	1,601	5.13	1.74
(%)	(1.3)	(2.0)	(5.4)	(14.9)	(24.1)	(24.2)	(16.1)	(7.0)	(3.1)	(1.9)	(100.0)				
9.0～	38	63	124	309	590	624	423	211	82	49	2,513	24	2,537	5.22	1.79
(%)	(1.5)	(2.5)	(4.9)	(12.3)	(23.5)	(24.8)	(16.8)	(8.4)	(3.3)	(1.9)	(100.0)				
合計	3,611	6,499	16,716	34,257	46,037	35,963	18,282	7,042	2,165	1,196	171,768	1,221	172,989	4.54	1.65
(%)	(2.1)	(3.8)	(9.7)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
記載なし	85	110	205	327	378	326	145	66	30	9	1,681	3,539	5,220	4.24	1.85
(%)	(5.1)	(6.5)	(12.2)	(19.5)	(22.5)	(19.4)	(8.6)	(3.9)	(1.8)	(0.5)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	4.53	4.64	4.88	5.09	5.26	5.39	5.49	5.59	5.64	5.68	5.22	5.14	5.22		
標準偏差	1.51	1.44	1.35	1.33	1.36	1.41	1.49	1.58	1.69	1.77	1.42	1.51	1.43		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査