

表 2707 体重減少率(%) 透析前ヘモグロビン濃度(g/dL)別 (施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前ヘモグロビン濃度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
5未満	6	5	14	44	59	46	20	6	3	2	205	5	210	4.54	1.66
(%)	(2.9)	(2.4)	(6.8)	(21.5)	(28.8)	(22.4)	(9.8)	(2.9)	(1.5)	(1.0)	(100.0)				
5～	6	20	24	42	54	47	32	14	4	1	244	7	251	4.50	1.82
(%)	(2.5)	(8.2)	(9.8)	(17.2)	(22.1)	(19.3)	(13.1)	(5.7)	(1.6)	(0.4)	(100.0)				
6～	32	39	72	109	146	100	75	20	9	5	607	10	617	4.28	1.91
(%)	(5.3)	(6.4)	(11.9)	(18.0)	(24.1)	(16.5)	(12.4)	(3.3)	(1.5)	(0.8)	(100.0)				
7～	104	170	279	462	555	392	257	119	40	32	2,410	35	2,445	4.34	1.94
(%)	(4.3)	(7.1)	(11.6)	(19.2)	(23.0)	(16.3)	(10.7)	(4.9)	(1.7)	(1.3)	(100.0)				
8～	266	447	883	1,575	2,046	1,632	944	393	124	96	8,406	91	8,497	4.49	1.84
(%)	(3.2)	(5.3)	(10.5)	(18.7)	(24.3)	(19.4)	(11.2)	(4.7)	(1.5)	(1.1)	(100.0)				
9～	611	1,151	2,913	5,508	7,250	5,874	3,165	1,261	414	256	28,403	205	28,608	4.56	1.70
(%)	(2.2)	(4.1)	(10.3)	(19.4)	(25.5)	(20.7)	(11.1)	(4.4)	(1.5)	(0.9)	(100.0)				
10～	1,089	2,089	5,576	11,978	16,163	12,643	6,349	2,408	737	384	59,416	353	59,769	4.56	1.62
(%)	(1.8)	(3.5)	(9.4)	(20.2)	(27.2)	(21.3)	(10.7)	(4.1)	(1.2)	(0.6)	(100.0)				
11～	978	1,765	4,684	9,766	13,156	10,168	4,911	1,842	561	280	48,111	304	48,415	4.51	1.61
(%)	(2.0)	(3.7)	(9.7)	(20.3)	(27.3)	(21.1)	(10.2)	(3.8)	(1.2)	(0.6)	(100.0)				
12～	390	585	1,630	3,428	4,696	3,612	1,799	688	184	87	17,099	109	17,208	4.52	1.62
(%)	(2.3)	(3.4)	(9.5)	(20.0)	(27.5)	(21.1)	(10.5)	(4.0)	(1.1)	(0.5)	(100.0)				
13～	99	140	412	793	1,183	875	447	182	57	29	4,217	43	4,260	4.56	1.67
(%)	(2.3)	(3.3)	(9.8)	(18.8)	(28.1)	(20.7)	(10.6)	(4.3)	(1.4)	(0.7)	(100.0)				
14～	39	67	157	327	405	319	143	51	16	14	1,538	13	1,551	4.44	1.69
(%)	(2.5)	(4.4)	(10.2)	(21.3)	(26.3)	(20.7)	(9.3)	(3.3)	(1.0)	(0.9)	(100.0)				
合計	3,620	6,478	16,644	34,032	45,713	35,708	18,142	6,984	2,149	1,186	170,656	1,175	171,831	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
記載なし	76	131	277	552	702	581	285	124	46	19	2,793	3,585	6,378	4.50	1.74
(%)	(2.7)	(4.7)	(9.9)	(19.8)	(25.1)	(20.8)	(10.2)	(4.4)	(1.6)	(0.7)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	10.61	10.58	10.68	10.73	10.75	10.74	10.67	10.65	10.59	10.47	10.71	10.54	10.71		
標準偏差	1.44	1.36	1.28	1.24	1.23	1.22	1.25	1.27	1.29	1.36	1.25	1.51	1.25		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査