

表 2711 体重減少率(%) 透析前収縮期血圧(mmHg)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前収縮期血圧	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	2	1	5	5	4	3	2		3		25	1	26	4.17	2.41
(%)	(8.0)	(4.0)	(20.0)	(20.0)	(16.0)	(12.0)	(8.0)		(12.0)		(100.0)				
60～	27	21	27	45	54	35	11	4	6		230	3	233	3.67	2.09
(%)	(11.7)	(9.1)	(11.7)	(19.6)	(23.5)	(15.2)	(4.8)	(1.7)	(2.6)		(100.0)				
80～	125	133	272	418	461	305	114	60	18	6	1,912	11	1,923	3.93	1.83
(%)	(6.5)	(7.0)	(14.2)	(21.9)	(24.1)	(16.0)	(6.0)	(3.1)	(0.9)	(0.3)	(100.0)				
100～	412	570	1,324	2,268	2,452	1,726	799	306	81	47	9,985	58	10,043	4.14	1.71
(%)	(4.1)	(5.7)	(13.3)	(22.7)	(24.6)	(17.3)	(8.0)	(3.1)	(0.8)	(0.5)	(100.0)				
120～	800	1,421	3,484	6,800	8,393	6,248	2,904	1,062	304	186	31,602	148	31,750	4.38	1.65
(%)	(2.5)	(4.5)	(11.0)	(21.5)	(26.6)	(19.8)	(9.2)	(3.4)	(1.0)	(0.6)	(100.0)				
140～	956	1,801	4,759	9,947	13,603	10,637	5,219	1,928	561	296	49,707	191	49,898	4.54	1.60
(%)	(1.9)	(3.6)	(9.6)	(20.0)	(27.4)	(21.4)	(10.5)	(3.9)	(1.1)	(0.6)	(100.0)				
160～	587	1,155	3,191	6,932	10,066	8,114	4,293	1,678	529	254	36,799	143	36,942	4.66	1.61
(%)	(1.6)	(3.1)	(8.7)	(18.8)	(27.4)	(22.0)	(11.7)	(4.6)	(1.4)	(0.7)	(100.0)				
180～	193	415	1,043	2,532	3,846	3,228	1,806	692	245	158	14,158	60	14,218	4.79	1.64
(%)	(1.4)	(2.9)	(7.4)	(17.9)	(27.2)	(22.8)	(12.8)	(4.9)	(1.7)	(1.1)	(100.0)				
200～	40	95	248	560	926	849	495	244	81	49	3,587	16	3,603	4.96	1.67
(%)	(1.1)	(2.6)	(6.9)	(15.6)	(25.8)	(23.7)	(13.8)	(6.8)	(2.3)	(1.4)	(100.0)				
220～	4	17	59	110	177	162	112	41	21	17	720	3	723	5.05	1.77
(%)	(0.6)	(2.4)	(8.2)	(15.3)	(24.6)	(22.5)	(15.6)	(5.7)	(2.9)	(2.4)	(100.0)				
合計	3,146	5,629	14,412	29,617	39,982	31,307	15,755	6,015	1,849	1,013	148,725	634	149,359	4.53	1.64
(%)	(2.1)	(3.8)	(9.7)	(19.9)	(26.9)	(21.1)	(10.6)	(4.0)	(1.2)	(0.7)	(100.0)				
記載なし	550	980	2,509	4,967	6,433	4,982	2,672	1,093	346	192	24,724	4,126	28,850	4.53	1.69
(%)	(2.2)	(4.0)	(10.1)	(20.1)	(26.0)	(20.2)	(10.8)	(4.4)	(1.4)	(0.8)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	142.96	146.66	148.01	150.12	152.28	153.67	155.39	156.02	157.38	158.57	151.91	149.22	151.90		
標準偏差	25.91	24.84	24.33	23.77	23.62	23.67	23.88	24.32	26.11	26.07	24.10	25.97	24.11		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査