

表 2712 体重減少率(%) 透析前拡張期血圧(mmHg)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前拡張 期血圧	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	1	4	4	9	14	12	3	4	1	1	53	2	55	4.67	2.14
(%)	(1.9)	(7.5)	(7.5)	(17.0)	(26.4)	(22.6)	(5.7)	(7.5)	(1.9)	(1.9)	(100.0)				
20～	28	33	68	123	107	83	52	17	4	2	517	7	524	4.09	1.81
(%)	(5.4)	(6.4)	(13.2)	(23.8)	(20.7)	(16.1)	(10.1)	(3.3)	(0.8)	(0.4)	(100.0)				
40～	441	661	1,591	2,657	3,282	2,227	1,009	422	126	64	12,480	75	12,555	4.23	1.72
(%)	(3.5)	(5.3)	(12.7)	(21.3)	(26.3)	(17.8)	(8.1)	(3.4)	(1.0)	(0.5)	(100.0)				
60～	1,531	2,819	7,157	14,371	18,580	13,968	6,752	2,383	707	390	68,658	283	68,941	4.44	1.62
(%)	(2.2)	(4.1)	(10.4)	(20.9)	(27.1)	(20.3)	(9.8)	(3.5)	(1.0)	(0.6)	(100.0)				
80～	654	1,267	3,237	7,125	9,952	7,993	3,982	1,576	461	239	36,486	144	36,630	4.60	1.61
(%)	(1.8)	(3.5)	(8.9)	(19.5)	(27.3)	(21.9)	(10.9)	(4.3)	(1.3)	(0.7)	(100.0)				
90～	328	576	1,629	3,596	5,302	4,493	2,427	970	308	154	19,783	80	19,863	4.72	1.63
(%)	(1.7)	(2.9)	(8.2)	(18.2)	(26.8)	(22.7)	(12.3)	(4.9)	(1.6)	(0.8)	(100.0)				
100～	138	232	649	1,585	2,477	2,302	1,377	581	217	133	9,691	40	9,731	4.94	1.68
(%)	(1.4)	(2.4)	(6.7)	(16.4)	(25.6)	(23.8)	(14.2)	(6.0)	(2.2)	(1.4)	(100.0)				
120～	9	14	34	79	145	134	104	36	14	24	593	1	594	5.21	1.81
(%)	(1.5)	(2.4)	(5.7)	(13.3)	(24.5)	(22.6)	(17.5)	(6.1)	(2.4)	(4.0)	(100.0)				
140～	1	1	1	6	10	10	4	7	2	1	43		43	5.41	1.94
(%)	(2.3)	(2.3)	(2.3)	(14.0)	(23.3)	(23.3)	(9.3)	(16.3)	(4.7)	(2.3)	(100.0)				
合計	3,131	5,607	14,370	29,551	39,869	31,222	15,710	5,996	1,840	1,008	148,304	632	148,936	4.53	1.64
(%)	(2.1)	(3.8)	(9.7)	(19.9)	(26.9)	(21.1)	(10.6)	(4.0)	(1.2)	(0.7)	(100.0)				
記載なし	565	1,002	2,551	5,033	6,546	5,067	2,717	1,112	355	197	25,145	4,128	29,273	4.53	1.69
(%)	(2.2)	(4.0)	(10.1)	(20.0)	(26.0)	(20.2)	(10.8)	(4.4)	(1.4)	(0.8)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	74.15	74.94	75.50	76.66	77.73	78.79	79.81	80.25	81.20	82.18	77.74	76.23	77.73		
標準偏差	15.01	14.29	14.14	14.12	14.21	14.35	14.76	15.24	15.62	17.01	14.47	15.85	14.47		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査