

表 2713 体重減少率(%) 透析前脈拍(拍/分)別(施設血液透析患者 透析回数週3回 透析歴2年以上)

透析前脈拍	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
40未満		4	17	17	27	24	15	8	1	1	114	1	115	4.72	1.65
(%)		(3.5)	(14.9)	(14.9)	(23.7)	(21.1)	(13.2)	(7.0)	(0.9)	(0.9)	(100.0)				
40～	25	49	134	278	383	314	157	63	20	13	1,436	2	1,438	4.62	1.64
(%)	(1.7)	(3.4)	(9.3)	(19.4)	(26.7)	(21.9)	(10.9)	(4.4)	(1.4)	(0.9)	(100.0)				
50～	220	418	1,120	2,371	3,354	2,510	1,255	452	146	65	11,911	51	11,962	4.54	1.59
(%)	(1.8)	(3.5)	(9.4)	(19.9)	(28.2)	(21.1)	(10.5)	(3.8)	(1.2)	(0.5)	(100.0)				
60～	697	1,354	3,600	7,541	10,150	8,138	4,049	1,535	472	254	37,790	148	37,938	4.56	1.63
(%)	(1.8)	(3.6)	(9.5)	(20.0)	(26.9)	(21.5)	(10.7)	(4.1)	(1.2)	(0.7)	(100.0)				
70～	892	1,659	4,389	8,934	12,180	9,671	4,883	1,856	560	294	45,318	183	45,501	4.55	1.63
(%)	(2.0)	(3.7)	(9.7)	(19.7)	(26.9)	(21.3)	(10.8)	(4.1)	(1.2)	(0.6)	(100.0)				
80～	680	1,121	2,781	5,771	7,638	5,864	3,001	1,150	336	217	28,559	105	28,664	4.51	1.67
(%)	(2.4)	(3.9)	(9.7)	(20.2)	(26.7)	(20.5)	(10.5)	(4.0)	(1.2)	(0.8)	(100.0)				
90～	314	490	1,150	2,501	3,336	2,494	1,249	492	159	98	12,283	48	12,331	4.50	1.67
(%)	(2.6)	(4.0)	(9.4)	(20.4)	(27.2)	(20.3)	(10.2)	(4.0)	(1.3)	(0.8)	(100.0)				
100～	114	181	376	734	990	789	423	147	59	25	3,838	20	3,858	4.48	1.73
(%)	(3.0)	(4.7)	(9.8)	(19.1)	(25.8)	(20.6)	(11.0)	(3.8)	(1.5)	(0.7)	(100.0)				
110～	34	37	75	180	244	159	87	39	15	8	878	3	881	4.49	1.84
(%)	(3.9)	(4.2)	(8.5)	(20.5)	(27.8)	(18.1)	(9.9)	(4.4)	(1.7)	(0.9)	(100.0)				
120～	14	16	47	67	102	80	38	23	5	8	400	5	405	4.57	1.91
(%)	(3.5)	(4.0)	(11.8)	(16.8)	(25.5)	(20.0)	(9.5)	(5.8)	(1.3)	(2.0)	(100.0)				
合計	2,990	5,329	13,689	28,394	38,404	30,043	15,157	5,765	1,773	983	142,527	566	143,093	4.54	1.64
(%)	(2.1)	(3.7)	(9.6)	(19.9)	(26.9)	(21.1)	(10.6)	(4.0)	(1.2)	(0.7)	(100.0)				
記載なし	706	1,280	3,232	6,190	8,011	6,246	3,270	1,343	422	222	30,922	4,194	35,116	4.51	1.68
(%)	(2.3)	(4.1)	(10.5)	(20.0)	(25.9)	(20.2)	(10.6)	(4.3)	(1.4)	(0.7)	(100.0)				
総計	3,696	6,609	16,921	34,584	46,415	36,289	18,427	7,108	2,195	1,205	173,449	4,760	178,209	4.53	1.65
(%)	(2.1)	(3.8)	(9.8)	(19.9)	(26.8)	(20.9)	(10.6)	(4.1)	(1.3)	(0.7)	(100.0)				
平均	76.44	75.30	74.64	74.59	74.45	74.31	74.45	74.64	74.79	75.71	74.56	75.24	74.56		
標準偏差	13.34	12.95	12.85	12.71	12.81	12.62	12.66	12.80	13.13	13.50	12.77	13.73	12.77		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査