

表 2717 体重減少率(%) 原疾患別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

原疾患	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	203	400	1,173	2,677	3,828	3,042	1,613	605	188	76	13,805	321	14,126	4.66	1.57
(%)	(1.5)	(2.9)	(8.5)	(19.4)	(27.7)	(22.0)	(11.7)	(4.4)	(1.4)	(0.6)	(100.0)				
慢性腎盂腎炎	9	16	33	68	109	75	50	22	3	4	389	9	398	4.63	1.69
(%)	(2.3)	(4.1)	(8.5)	(17.5)	(28.0)	(19.3)	(12.9)	(5.7)	(0.8)	(1.0)	(100.0)				
急速進行性糸球体腎炎	4	10	13	37	62	64	36	10	5	4	245	5	250	4.89	1.67
(%)	(1.6)	(4.1)	(5.3)	(15.1)	(25.3)	(26.1)	(14.7)	(4.1)	(2.0)	(1.6)	(100.0)				
妊娠腎/妊娠中毒症	3	9	24	67	93	86	43	14	3		342	6	348	4.65	1.46
(%)	(0.9)	(2.6)	(7.0)	(19.6)	(27.2)	(25.1)	(12.6)	(4.1)	(0.9)		(100.0)				
その他の分類不能の腎炎	4	2	10	35	53	40	17	14	7		182	4	186	4.86	1.62
(%)	(2.2)	(1.1)	(5.5)	(19.2)	(29.1)	(22.0)	(9.3)	(7.7)	(3.8)		(100.0)				
多発性嚢胞腎	24	46	150	288	382	251	109	39	4	3	1,296	34	1,330	4.34	1.53
(%)	(1.9)	(3.5)	(11.6)	(22.2)	(29.5)	(19.4)	(8.4)	(3.0)	(0.3)	(0.2)	(100.0)				
腎硬化症	40	84	231	471	597	440	227	94	24	14	2,222	30	2,252	4.48	1.61
(%)	(1.8)	(3.8)	(10.4)	(21.2)	(26.9)	(19.8)	(10.2)	(4.2)	(1.1)	(0.6)	(100.0)				
悪性高血圧	7	3	18	40	90	55	35	13	3	1	265	5	270	4.75	1.54
(%)	(2.6)	(1.1)	(6.8)	(15.1)	(34.0)	(20.8)	(13.2)	(4.9)	(1.1)	(0.4)	(100.0)				
糖尿病性腎症	190	352	969	2,236	2,914	2,303	1,066	371	118	57	10,576	269	10,845	4.53	1.56
(%)	(1.8)	(3.3)	(9.2)	(21.1)	(27.6)	(21.8)	(10.1)	(3.5)	(1.1)	(0.5)	(100.0)				
SLE腎炎	5	18	24	57	72	80	40	27	6	2	331	6	337	4.81	1.74
(%)	(1.5)	(5.4)	(7.3)	(17.2)	(21.8)	(24.2)	(12.1)	(8.2)	(1.8)	(0.6)	(100.0)				
アミロイド腎		3	1	8	12	13	3	2		1	43		43	4.77	1.55
(%)		(7.0)	(2.3)	(18.6)	(27.9)	(30.2)	(7.0)	(4.7)		(2.3)	(100.0)				
痛風腎	4	2	16	27	42	26	15	4			136	6	142	4.38	1.46
(%)	(2.9)	(1.5)	(11.8)	(19.9)	(30.9)	(19.1)	(11.0)	(2.9)			(100.0)				
先天性代謝異常による腎不全		1	5	4	8	10	5	2	3	1	39	2	41	5.17	1.87
(%)		(2.6)	(12.8)	(10.3)	(20.5)	(25.6)	(12.8)	(5.1)	(7.7)	(2.6)	(100.0)				
腎・尿路結核	1		3	5	2	7	2				20		20	4.23	1.43
(%)	(5.0)		(15.0)	(25.0)	(10.0)	(35.0)	(10.0)				(100.0)				
腎・尿路結石	1	3	3	13	20	15	5	3			63	1	64	4.51	1.42
(%)	(1.6)	(4.8)	(4.8)	(20.6)	(31.7)	(23.8)	(7.9)	(4.8)			(100.0)				
腎・尿路腫瘍	3	1	5	12	20	15	4		2		62		62	4.42	1.56
(%)	(4.8)	(1.6)	(8.1)	(19.4)	(32.3)	(24.2)	(6.5)		(3.2)		(100.0)				
閉塞性尿路障害	1	1	6	16	16	8	12	5		1	66	3	69	4.79	1.75
(%)	(1.5)	(1.5)	(9.1)	(24.2)	(24.2)	(12.1)	(18.2)	(7.6)		(1.5)	(100.0)				
骨髄腫		1	2	3	1	4					11		11	3.95	1.45
(%)		(9.1)	(18.2)	(27.3)	(9.1)	(36.4)					(100.0)				
腎形成不全	1	3	10	9	29	25	8	8	1		94	1	95	4.81	1.53
(%)	(1.1)	(3.2)	(10.6)	(9.6)	(30.9)	(26.6)	(8.5)	(8.5)	(1.1)		(100.0)				
不明	67	108	253	598	744	603	311	115	47	16	2,862	92	2,954	4.54	1.67
(%)	(2.3)	(3.8)	(8.8)	(20.9)	(26.0)	(21.1)	(10.9)	(4.0)	(1.6)	(0.6)	(100.0)				
再導入	1	3	24	80	88	96	50	22	8	5	377	8	385	4.96	1.55
(%)	(0.3)	(0.8)	(6.4)	(21.2)	(23.3)	(25.5)	(13.3)	(5.8)	(2.1)	(1.3)	(100.0)				
その他	19	25	57	142	195	166	74	38	14	5	735	8	743	4.64	1.70
(%)	(2.6)	(3.4)	(7.8)	(19.3)	(26.5)	(22.6)	(10.1)	(5.2)	(1.9)	(0.7)	(100.0)				
合計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
記載なし															
(%)															
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査