

表 2722 体重減少率(%) Body Mass Index別(血液透析濾過患者 透析回数週3回 透析歴2年以上)

Body Mass Index	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
14未満	13	6	9	21	15	23	22	10	3	3	125	13	138	4.60	2.35
(%)	(10.4)	(4.8)	(7.2)	(16.8)	(12.0)	(18.4)	(17.6)	(8.0)	(2.4)	(2.4)	(100.0)				
14～	27	36	76	137	190	209	133	94	34	25	961	3	964	5.04	2.02
(%)	(2.8)	(3.7)	(7.9)	(14.3)	(19.8)	(21.7)	(13.8)	(9.8)	(3.5)	(2.6)	(100.0)				
16～	70	113	247	519	827	851	591	283	113	39	3,653	6	3,659	5.02	1.77
(%)	(1.9)	(3.1)	(6.8)	(14.2)	(22.6)	(23.3)	(16.2)	(7.7)	(3.1)	(1.1)	(100.0)				
18～	102	198	545	1,169	1,664	1,657	931	398	123	55	6,842	7	6,849	4.83	1.65
(%)	(1.5)	(2.9)	(8.0)	(17.1)	(24.3)	(24.2)	(13.6)	(5.8)	(1.8)	(0.8)	(100.0)				
20～	136	215	648	1,404	2,086	1,689	850	278	86	35	7,427	5	7,432	4.60	1.58
(%)	(1.8)	(2.9)	(8.7)	(18.9)	(28.1)	(22.7)	(11.4)	(3.7)	(1.2)	(0.5)	(100.0)				
22～	97	191	560	1,284	1,719	1,177	539	141	33	20	5,761	1	5,762	4.42	1.47
(%)	(1.7)	(3.3)	(9.7)	(22.3)	(29.8)	(20.4)	(9.4)	(2.4)	(0.6)	(0.3)	(100.0)				
24～	61	117	363	880	1,139	661	250	77	14		3,562		3,562	4.29	1.37
(%)	(1.7)	(3.3)	(10.2)	(24.7)	(32.0)	(18.6)	(7.0)	(2.2)	(0.4)		(100.0)				
26～	20	78	181	490	574	359	121	29	2	1	1,855		1,855	4.24	1.30
(%)	(1.1)	(4.2)	(9.8)	(26.4)	(30.9)	(19.4)	(6.5)	(1.6)	(0.1)	(0.1)	(100.0)				
28～	10	30	113	287	318	184	49	7			998		998	4.15	1.22
(%)	(1.0)	(3.0)	(11.3)	(28.8)	(31.9)	(18.4)	(4.9)	(0.7)			(100.0)				
30～	12	28	102	235	270	153	24	7	1	2	834	2	836	4.10	1.25
(%)	(1.4)	(3.4)	(12.2)	(28.2)	(32.4)	(18.3)	(2.9)	(0.8)	(0.1)	(0.2)	(100.0)				
合計	548	1,012	2,844	6,426	8,802	6,963	3,510	1,324	409	180	32,018	37	32,055	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.1)	(27.5)	(21.7)	(11.0)	(4.1)	(1.3)	(0.6)	(100.0)				
記載なし	39	79	186	467	575	461	215	84	27	10	2,143	773	2,916	4.55	1.58
(%)	(1.8)	(3.7)	(8.7)	(21.8)	(26.8)	(21.5)	(10.0)	(3.9)	(1.3)	(0.5)	(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	21.02	21.73	22.07	22.33	22.02	21.30	20.48	19.72	19.14	18.97	21.58	15.78	21.58		
標準偏差	3.63	3.72	3.70	3.74	3.59	3.49	3.13	2.94	2.50	2.73	3.61	6.33	3.62		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査