

表 2723 体重減少率(%) Body Mass Index別 (血液透析濾過患者 男性のみ 透析回数週3回 透析歴2年以上)

Body Mass Index	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
14未満	3	4	3	8	3	6	4	2			33	7	40	3.92	2.10
(%)	(9.1)	(12.1)	(9.1)	(24.2)	(9.1)	(18.2)	(12.1)	(6.1)			(100.0)				
14～	9	17	29	42	66	63	45	36	11	5	323	1	324	4.94	1.97
(%)	(2.8)	(5.3)	(9.0)	(13.0)	(20.4)	(19.5)	(13.9)	(11.1)	(3.4)	(1.5)	(100.0)				
16～	26	55	117	238	384	394	249	123	51	20	1,657	3	1,660	5.00	1.77
(%)	(1.6)	(3.3)	(7.1)	(14.4)	(23.2)	(23.8)	(15.0)	(7.4)	(3.1)	(1.2)	(100.0)				
18～	62	111	312	640	965	919	537	242	79	38	3,905	3	3,908	4.86	1.68
(%)	(1.6)	(2.8)	(8.0)	(16.4)	(24.7)	(23.5)	(13.8)	(6.2)	(2.0)	(1.0)	(100.0)				
20～	92	134	439	938	1,385	1,071	571	190	62	26	4,908	4	4,912	4.61	1.59
(%)	(1.9)	(2.7)	(8.9)	(19.1)	(28.2)	(21.8)	(11.6)	(3.9)	(1.3)	(0.5)	(100.0)				
22～	72	135	397	865	1,207	841	392	112	25	12	4,058	1	4,059	4.44	1.48
(%)	(1.8)	(3.3)	(9.8)	(21.3)	(29.7)	(20.7)	(9.7)	(2.8)	(0.6)	(0.3)	(100.0)				
24～	42	80	247	615	796	476	190	56	10		2,512		2,512	4.31	1.38
(%)	(1.7)	(3.2)	(9.8)	(24.5)	(31.7)	(18.9)	(7.6)	(2.2)	(0.4)		(100.0)				
26～	16	47	118	337	408	269	94	23	2	1	1,315		1,315	4.30	1.32
(%)	(1.2)	(3.6)	(9.0)	(25.6)	(31.0)	(20.5)	(7.1)	(1.7)	(0.2)	(0.1)	(100.0)				
28～	8	18	67	192	218	142	36	3			684		684	4.20	1.22
(%)	(1.2)	(2.6)	(9.8)	(28.1)	(31.9)	(20.8)	(5.3)	(0.4)			(100.0)				
30～	8	15	54	152	184	117	15	4	1		550	2	552	4.19	1.19
(%)	(1.5)	(2.7)	(9.8)	(27.6)	(33.5)	(21.3)	(2.7)	(0.7)	(0.2)		(100.0)				
合計	338	616	1,783	4,027	5,616	4,298	2,133	791	241	102	19,945	21	19,966	4.58	1.57
(%)	(1.7)	(3.1)	(8.9)	(20.2)	(28.2)	(21.5)	(10.7)	(4.0)	(1.2)	(0.5)	(100.0)				
記載なし	25	39	106	279	374	289	132	49	18	5	1,316	494	1,810	4.59	1.55
(%)	(1.9)	(3.0)	(8.1)	(21.2)	(28.4)	(22.0)	(10.0)	(3.7)	(1.4)	(0.4)	(100.0)				
総計	363	655	1,889	4,306	5,990	4,587	2,265	840	259	107	21,261	515	21,776	4.58	1.57
(%)	(1.7)	(3.1)	(8.9)	(20.3)	(28.2)	(21.6)	(10.7)	(4.0)	(1.2)	(0.5)	(100.0)				
平均	21.73	21.93	22.29	22.71	22.40	21.87	21.09	20.27	19.75	19.49	22.04	17.12	22.04		
標準偏差	3.35	3.57	3.43	3.58	3.46	3.48	3.03	2.83	2.44	2.07	3.47	6.73	3.48		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査