

表 2725 体重減少率(%) 透析前BUN(mg/dL)別(血液透析濾過患者 透析回数週3回 透析歴2年以上)

透析前BUN	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	12	5	5	6	4	3	2				37	2	39	2.46	2.01
(%)	(32.4)	(13.5)	(13.5)	(16.2)	(10.8)	(8.1)	(5.4)				(100.0)				
20～	35	46	59	71	40	27	12	7	2	2	301	8	309	3.22	1.90
(%)	(11.6)	(15.3)	(19.6)	(23.6)	(13.3)	(9.0)	(4.0)	(2.3)	(0.7)	(0.7)	(100.0)				
30～	80	132	264	414	355	230	103	38	16	10	1,642	11	1,653	3.88	1.74
(%)	(4.9)	(8.0)	(16.1)	(25.2)	(21.6)	(14.0)	(6.3)	(2.3)	(1.0)	(0.6)	(100.0)				
40～	127	238	617	1,147	1,199	889	362	123	34	14	4,750	35	4,785	4.20	1.60
(%)	(2.7)	(5.0)	(13.0)	(24.1)	(25.2)	(18.7)	(7.6)	(2.6)	(0.7)	(0.3)	(100.0)				
50～	117	293	812	1,846	2,302	1,676	796	293	93	35	8,263	55	8,318	4.47	1.54
(%)	(1.4)	(3.5)	(9.8)	(22.3)	(27.9)	(20.3)	(9.6)	(3.5)	(1.1)	(0.4)	(100.0)				
60～	101	200	698	1,724	2,591	1,977	957	371	85	53	8,757	47	8,804	4.67	1.50
(%)	(1.2)	(2.3)	(8.0)	(19.7)	(29.6)	(22.6)	(10.9)	(4.2)	(1.0)	(0.6)	(100.0)				
70～	69	113	345	1,044	1,756	1,524	767	300	105	30	6,053	33	6,086	4.86	1.50
(%)	(1.1)	(1.9)	(5.7)	(17.2)	(29.0)	(25.2)	(12.7)	(5.0)	(1.7)	(0.5)	(100.0)				
80～	30	35	139	436	766	706	463	164	57	20	2,816	15	2,831	5.04	1.51
(%)	(1.1)	(1.2)	(4.9)	(15.5)	(27.2)	(25.1)	(16.4)	(5.8)	(2.0)	(0.7)	(100.0)				
90～	8	11	50	132	248	278	186	81	32	20	1,046	5	1,051	5.28	1.60
(%)	(0.8)	(1.1)	(4.8)	(12.6)	(23.7)	(26.6)	(17.8)	(7.7)	(3.1)	(1.9)	(100.0)				
100～	3	1	15	33	65	75	46	24	8	6	276	3	279	5.33	1.61
(%)	(1.1)	(0.4)	(5.4)	(12.0)	(23.6)	(27.2)	(16.7)	(8.7)	(2.9)	(2.2)	(100.0)				
110～	1		1	7	20	18	21	4	2		74		74	5.39	1.44
(%)	(1.4)		(1.4)	(9.5)	(27.0)	(24.3)	(28.4)	(5.4)	(2.7)		(100.0)				
120～			2	3	7	3	3	1	1		20		20	4.98	1.49
(%)			(10.0)	(15.0)	(35.0)	(15.0)	(15.0)	(5.0)	(5.0)		(100.0)				
合計	583	1,074	3,007	6,863	9,353	7,406	3,718	1,406	435	190	34,035	214	34,249	4.59	1.58
(%)	(1.7)	(3.2)	(8.8)	(20.2)	(27.5)	(21.8)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
記載なし	4	17	23	30	24	18	7	2	1		126	596	722	3.72	1.62
(%)	(3.2)	(13.5)	(18.3)	(23.8)	(19.0)	(14.3)	(5.6)	(1.6)	(0.8)		(100.0)				
総計	587	1,091	3,030	6,893	9,377	7,424	3,725	1,408	436	190	34,161	810	34,971	4.59	1.59
(%)	(1.7)	(3.2)	(8.9)	(20.2)	(27.4)	(21.7)	(10.9)	(4.1)	(1.3)	(0.6)	(100.0)				
平均	53.54	53.73	57.00	59.86	62.86	64.42	66.23	66.77	68.14	67.48	62.25	59.15	62.23		
標準偏差	17.50	14.76	14.74	14.37	14.29	14.62	15.31	15.24	16.31	17.01	15.07	16.67	15.08		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査