

表 2737 体重減少率(%) %クレアチニン産生速度(%)別(血液透析濾過患者 男性のみ 透析回数週3回 透析歴2年以上)

%クレアチニン 産生速度	1.0未満	1.0～	2.0～	3.0～	4.0～	5.0～	6.0～	7.0～	8.0～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	76	102	149	228	199	146	59	33	9	1	1,002		1,002	3.73	1.82
(%)	(7.6)	(10.2)	(14.9)	(22.8)	(19.9)	(14.6)	(5.9)	(3.3)	(0.9)	(0.1)	(100.0)				
60～	48	64	108	197	208	132	75	38	10	6	886		886	4.11	1.84
(%)	(5.4)	(7.2)	(12.2)	(22.2)	(23.5)	(14.9)	(8.5)	(4.3)	(1.1)	(0.7)	(100.0)				
70～	48	76	181	360	400	275	143	56	22	9	1,570	1	1,571	4.31	1.70
(%)	(3.1)	(4.8)	(11.5)	(22.9)	(25.5)	(17.5)	(9.1)	(3.6)	(1.4)	(0.6)	(100.0)				
80～	51	86	251	559	670	507	254	100	37	14	2,529		2,529	4.50	1.63
(%)	(2.0)	(3.4)	(9.9)	(22.1)	(26.5)	(20.0)	(10.0)	(4.0)	(1.5)	(0.6)	(100.0)				
90～	37	81	309	682	845	668	393	136	39	17	3,207	1	3,208	4.63	1.54
(%)	(1.2)	(2.5)	(9.6)	(21.3)	(26.3)	(20.8)	(12.3)	(4.2)	(1.2)	(0.5)	(100.0)				
100～	33	77	294	704	1,053	822	382	142	57	27	3,591	1	3,592	4.70	1.51
(%)	(0.9)	(2.1)	(8.2)	(19.6)	(29.3)	(22.9)	(10.6)	(4.0)	(1.6)	(0.8)	(100.0)				
110～	27	53	230	607	1,043	809	394	133	40	15	3,351	1	3,352	4.77	1.44
(%)	(0.8)	(1.6)	(6.9)	(18.1)	(31.1)	(24.1)	(11.8)	(4.0)	(1.2)	(0.4)	(100.0)				
120～	18	40	151	421	707	569	286	98	23	7	2,320	1	2,321	4.77	1.39
(%)	(0.8)	(1.7)	(6.5)	(18.1)	(30.5)	(24.5)	(12.3)	(4.2)	(1.0)	(0.3)	(100.0)				
130～	6	23	93	255	388	302	131	50	8	4	1,260	3	1,263	4.71	1.42
(%)	(0.5)	(1.8)	(7.4)	(20.2)	(30.8)	(24.0)	(10.4)	(4.0)	(0.6)	(0.3)	(100.0)				
140～	1	13	44	161	278	222	76	24	9	4	832	8	840	4.76	1.31
(%)	(0.1)	(1.6)	(5.3)	(19.4)	(33.4)	(26.7)	(9.1)	(2.9)	(1.1)	(0.5)	(100.0)				
合計	345	615	1,810	4,174	5,791	4,452	2,193	810	254	104	20,548	16	20,564	4.58	1.56
(%)	(1.7)	(3.0)	(8.8)	(20.3)	(28.2)	(21.7)	(10.7)	(3.9)	(1.2)	(0.5)	(100.0)				
記載なし	18	40	79	132	199	135	72	30	5	3	713	499	1,212	4.38	1.67
(%)	(2.5)	(5.6)	(11.1)	(18.5)	(27.9)	(18.9)	(10.1)	(4.2)	(0.7)	(0.4)	(100.0)				
総計	363	655	1,889	4,306	5,990	4,587	2,265	840	259	107	21,261	515	21,776	4.58	1.57
(%)	(1.7)	(3.1)	(8.9)	(20.3)	(28.2)	(21.6)	(10.7)	(4.0)	(1.2)	(0.5)	(100.0)				
平均	79.64	86.83	95.63	99.98	104.21	105.12	103.76	101.78	100.74	100.46	101.65	146.17	101.69		
標準偏差	27.59	28.08	24.89	23.88	23.03	22.55	21.08	23.13	21.94	21.50	23.80	50.82	23.86		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査